

Flows, View Obstructions, and the Lonely Runner

Wojciech Bienia

ENSIMAG, Laboratoire Leibniz-IMAG, France

Luis Goddyn* and Pavol Gvozďjak

Simon Fraser University, Vancouver, Canada

András Sebő†

CNRS, Laboratoire Leibniz-IMAG, France

and

Michael Tarsi‡

Tel Aviv University, Israel

Received August 29, 1996

We prove the following result: Let G be an undirected graph. If G has a nowhere zero flow with at most k different values, then it also has one with values from the set $\{1, \dots, k\}$. When $k \geq 5$, this is a trivial consequence of Seymour's "six-flow theorem". When $k \leq 4$ our proof is based on a lovely number theoretic problem which we call the "Lonely Runner Conjecture:" Suppose k runners having nonzero constant speeds run laps on a unit-length circular track. Then there is a time at which all runners are at least $1/(k+1)$ from their common starting point. This conjecture appears to have been formulated by J. Wills (*Monatsch. Math.* **71**, 1967) and independently by T. Cusick (*Aequationes Math.* **9**, 1973). This conjecture has been verified for $k \leq 4$ by Cusick and Pomerance (*J. Number Theory* **19**, 1984) in a complicated argument involving exponential sums and electronic case checking. A major part of this paper is an elementary self-contained proof of the case $k = 4$ of the Lonely Runner Conjecture. © 1998 Academic Press

* Research supported by NSERC of Canada. E-mail: goddyn@math.sfu.ca.

† Research supported by W. Cunningham's grant from NSERC of Canada. E-mail: sebo@imag.fr.

‡ Research supported in part by a grant from the Israel Science Foundation. E-mail: tarsi@math.tau.ac.il.

1. INTRODUCTION

Let $G = (V, E)$ be an undirected graph. A *nowhere zero flow* of G is an orientation of G supplied with a vector $f = (f_e)$ of positive integers indexed by $E(G)$, such that for every $v \in V(G)$ the sum of f_e on edges entering v is the same as that on edges leaving v . The number f_e is called the *value* of the edge e . The theory of nowhere zero flows is a major topic in combinatorics related to graph coloring and the cycle double cover conjecture; see [9, 14, 16]. The main result of this paper is the following.

THEOREM 1.1. *Let G be an undirected graph. If G has a nowhere zero flow with at most k distinct values, then it also has one with all values from the set $\{1, \dots, k\}$.*

In view of the matroid duality [16, 15, 9, 11, 14] between vertex colorings and nowhere zero flows there is a cographic analogue to Theorem 1.1. A *coloring* of G is a function $c: V(G) \rightarrow \mathbb{R}$, so that for all $xy \in E$, $c(x) \neq c(y)$.

THEOREM 1.2. *If G has a coloring with real numbers so that the set $\{|c(x) - c(y)|: xy \in E\}$ has at most k distinct values, then G has a $(k + 1)$ -coloring (and thus one where $|c(x) - c(y)| \in \{1, \dots, k\}$ for all $xy \in E$.)*

Theorem 1.2 is easy to prove. By orienting each edge toward the endpoint with the larger color and identifying the color classes, one obtains an acyclic digraph having maximum out-degree k . An easy greedy algorithm results in a $(k + 1)$ -coloring of G .

Theorem 1.1 is more difficult. Our proof relies on Seymour's *six-flow theorem* [13] and a number theoretic result of Cusick and Pomerance [6] to which we give a short proof. We state here the six-flow theorem. A graph is called *bridgeless*, if it has no bridge, where $e \in E$ is a *bridge* if $G - e$ has more components than G .

THEOREM 1.3. *Every bridgeless graph has a nowhere zero flow with values from the set $\{1, \dots, 5\}$.*

There is a common generalization of Theorems 1.1 and 1.2 regarding flows in regular matroids (see [11, 15]) which is strongly suggested by Seymour's *regular matroid decomposition theorem* [12]. A matrix is *totally unimodular* if every subdeterminant belongs to $\{0, \pm 1\}$.

Conjecture 1.4. Let A be a totally unimodular matrix and suppose that $Af = 0$ has a real solution $f = (f_e)$, where each f_e is nonzero and where $|\{f_e: e \in E(G)\}| \leq k$. Then there exists a solution $f' = (f'_e)$ with each $f'_e \in \{1, 2, \dots, k\}$.

The analogous statement concerning *group-valued flows* [16, 9] is false. For example, the graph with two vertices and three parallel edges has a flow with range $\{1\}$ in Z_3 , but not in the integers.

The paper is organized as follows. In Section 2, Conjecture 1.4 is reduced to the “Lonely Runner Problem”; in particular, Theorem 1.1 is reduced to the special case $k \leq 4$. A general proof technique for this problem is introduced in Section 3 and is applied to the case $k = 4$ in Section 4.

2. RUNNERS AND FLOWS

Let us informally state the *Lonely Runner Problem*: At time zero, k participants depart from the origin of a unit length circular track to run repeated laps. Each runner maintains a constant nonzero speed. Is it true that regardless of what the speeds are, there exists a time at which the k runners are simultaneously at least $1/(k + 1)$ units from the starting point? The term “lonely runner” reflects an equivalent formulation in which there are $k + 1$ runners with distinct speeds. Is there a time at which a given runner is “lonely,” that is, at distance at least $1/(k + 1)$ from the others? This poetic title (given by the second author) made its way through an internet inquiry (of the second and last authors) up to the cover page of a public relations booklet for the Weizmann Institute in Israel [22].

We introduce some notation. The sets of real numbers and positive integers are denoted \mathbb{R} and \mathbb{N} , respectively. The residue class of $a \in \mathbb{R}$ modulo 1 (called the *fractional part* of a) is denoted by $\langle a \rangle$. We view the unit-length circle C as the set $\{\langle a \rangle : a \in \mathbb{R}\}$, which we frequently identify with the real interval $[0, 1)$. An instance of the Lonely Runner Problem consists of a set of *runners* $R := \{1, 2, \dots, k\}$ and a *speed vector* $v := (v_1, \dots, v_k)$ having nonzero real entries. At time $t = 0$, each $r \in R$ begins running on C from the point 0 maintaining the constant speed v_r . The *position* of runner r on C at time t is $\langle tv_r \rangle$. The position of R at time t is the vector $\langle tv \rangle := (\langle tv_1 \rangle, \dots, \langle tv_k \rangle) \in [0, 1)^k$. A vector $x = (x_1, \dots, x_k) \in [0, 1)^k$ is a *position* (for the speed vector v) if there exists $t \in \mathbb{R}$ with $x = \langle tv \rangle$. The set of all positions is denoted $X = X(v) \subseteq [0, 1)^k$. The distance between two points on C is the length of the shorter of the two (arc) intervals between them. We say that $r \in R$ is *distant* (from 0) in $x \in X$ or *at time* t if $x_r = \langle tv_r \rangle \in [1/(k + 1), k/(k + 1)]$. A subset $R' \subseteq R$ is *distant* (in some position x) if each $r \in R'$ is distant in x . (Here, k is understood to equal $|R|$, not $|R'|$).

The aforementioned internet inquiry led us to the following assertion, which we call the *Lonely Runner Conjecture*. This conjecture appears to have been introduced by Wills [17] and again, independently by Cusick [3].

Conjecture 2.1. For all $k \in \mathbb{N}$ and $v \in (\mathbb{R} - \{0\})^k$, there exists a position where R is distant.

This problem appears in two different contexts. Cusick [3–6] was motivated by a beautiful application in n dimensional geometry—*view obstruction problems*. Our statement of the problem is closer to the diophantine approximation approach of Wills [1, 17–21]. A more general conjecture appears in [2]. The cases $k=2, 3, 4$ were first proved in [17, 1, 6], respectively.

THEOREM 2.2. *If $k \leq 4$, then for any $v \in (\mathbb{R} - \{0\})^k$ there exists a time at which R is distant.*

The proof by Cusick and Pomerance [6] of the case $k=4$ is not easy and requires a computer check. In Sections 3 and 4 we provide a simple self-contained proof. Section 3 also contains a very short proof for the case $k=3$. We now prove Theorem 1.1 using Theorems 2.2 and 1.3.

Proof of Theorem 1.1. Let f be a nowhere zero flow with k different values. If $k \geq 5$, then the result is a trivial consequence of Theorem 1.3 since any graph having a nowhere zero flow must be bridgeless. If $k \leq 4$, then by Theorem 2.2 there exists $t \in \mathbb{R}$ such that the fractional part of each entry of tf is in the interval $[1/(k+1), k/(k+1)]$. The flow tf is a feasible flow in the edge-capacitated network (G, l, u) , where $l = \lfloor tf \rfloor$ and $u = \lceil tf \rceil$ (we take floors and ceilings componentwise). But then there also exists a feasible integer-valued flow for (G, l, u) , (Ford and Fulkerson, [7]), in which each edge e has value either $\lfloor tf_e \rfloor$ or $\lceil tf_e \rceil$. Let us denote this flow by $\lfloor tf \rfloor$. Thus $tf - \lfloor tf \rfloor$ is a flow with all entries in $[-k/(k+1), -1/(k+1)] \cup [1/(k+1), k/(k+1)]$. Multiplying this flow by $k+1$ and reorienting the edges corresponding to negative entries yields a flow with values in $[1, k]$. Again, there also exists then an integer flow with values in $[1, k]$. ■

Note. We may loosely denote the final flow in the proof of Theorem 1.1 as $\lfloor (k+1)(f - \lfloor tf \rfloor) \rfloor$.

We remark that this proof can be directly generalized to flows in regular matroids by applying Hoffman's theorem [8] in order to define $f' = \lfloor (k+1)(f - \lfloor tf \rfloor) \rfloor$. Thus, Conjecture 1.4 is a weak form of the Lonely Runner Conjecture.

THEOREM 2.3. *For any $k \in \mathbb{N}$, if the Lonely Runner Conjecture holds true for k runners, then the statement of Conjecture 1.4 holds true for that particular value of k .*

The remainder of this paper is devoted to the Lonely Runner Conjecture. Wills [17] reduced the Lonely Runner Conjecture from the case of irrational

speeds to the rational case. So when proving any case $k \geq 1$, one can assume without loss of generality that $v \in \mathbb{N}^k$, whence the speeds express the number of laps the runners make in unit time. One can further assume that $t \in [0, 1)$, although there is usually no advantage in doing so.

Proof of Theorem 2.2 when $k \leq 2$. The case $k = 1$ is trivial. In case $k = 2$ we prove a stronger statement:

Suppose $v_1, v_2 \in \mathbb{N}$ are relatively prime speeds. At any time t , the nearer runner has distance at most $\lfloor (v_1 + v_2)/2 \rfloor / (v_1 + v_2)$. Moreover, this bound is achieved at time $t = \tau / (v_1 + v_2)$ for some $\tau \in \mathbb{N}$.

Whenever the distance from 0 to the nearer runner is maximum, we have $\langle tv_1 \rangle = 1 - \langle tv_2 \rangle$. This equality holds if and only if t is an integer multiple of $1/(v_1 + v_2)$. For such t , both runners are at distance $a/(v_1 + v_2)$ for some integer $a \leq \lfloor (v_1 + v_2)/2 \rfloor$. Since $\gcd(v_1, v_1 + v_2) = 1$ we can solve the congruence $v_1 \tau \equiv \lfloor (v_1 + v_2)/2 \rfloor \pmod{v_1 + v_2}$, to obtain a time at which the bound on a is achieved, proving the statement. ■

3. PREJUMPS

We state the fact that *the set X of positions is closed under addition modulo 1* in a particular form suggesting a technique used by all the proofs hereafter.

(1) If $x_1, x_2 \in X$ and $\alpha \in \mathbb{Z}$, then the vector $x = \langle x_1 + \alpha x_2 \rangle \in [0, 1)^k$ is also in X . If moreover, $x_1 = \langle t_1 v \rangle$, $x_2 = \langle t_2 v \rangle$, and $t \equiv t_1 + \alpha t_2 \pmod{1}$, then $x = \langle tv \rangle$.

Our use of (1) is as follows. We first note the existence of certain “key” positions in X which we call *prejumps*. In the proof of our main result, it sometimes becomes convenient to add one of these prejumps to a position that has already been constructed, thereby obtaining a position in which all runners are distant. Our first example of prejumps will be used in a short proof of the case $k = 3$. (Compare with the proofs in [1, 3].)

(2) Let $v \in \mathbb{N}^k$, $k \geq 3$. If $\gcd(v_1, \dots, v_{k-1})$ does not divide v_k , then there exists a time when R is distant if and only if there exists a time when $R \setminus \{k\}$ is distant.

Proof. Let $d \geq 2$ be the greatest common divisor defined in the statement, and suppose without loss of generality that $\gcd(d, v_k) = 1$. Then

$$\left\langle \frac{0}{d} v_r \right\rangle = \left\langle \frac{1}{d} v_r \right\rangle = \dots = \left\langle \frac{d-1}{d} v_r \right\rangle = 0 \quad \text{for } r = 1, \dots, k-1,$$

whereas

$$\left\{ \left\langle \frac{0}{k} v_k \right\rangle, \left\langle \frac{1}{d} v_k \right\rangle, \dots, \left\langle \frac{d-1}{d} v_k \right\rangle \right\} = \left\{ \frac{0}{d}, \frac{1}{d}, \dots, \frac{d-1}{d} \right\}.$$

Let now $x = \langle tv \rangle$ be a position where $R \setminus \{k\}$ is distant. Since $R \setminus \{k\}$ is also distant in each of the d positions $\langle x + (j/d)v \rangle$ ($1 = 0, 1, \dots, d-1$), it suffices to show that k is distant in one of these positions. However, this follows from the fact that $1/d$ is at most the length $1 - 2/(k+1)$ of the interval of distant positions since $k \geq 3$ and $d \geq 2$.

Proof of Theorem 2.2 when $k \leq 3$. We assume that the speeds v_1, v_2, v_3 are distinct positive integers having no common factor. If all three speeds are odd, then $\langle \frac{1}{2}v \rangle = (\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$, so we may assume that v_2 is even. By (2) we may further assume that v_1 and v_3 are odd. So $\langle \frac{1}{2}v \rangle = (\frac{1}{2}, 0, \frac{1}{2})$, and this will provide our prejump $x_1 = \langle t_1 v \rangle$, $t_1 := \frac{1}{2}$.

Consider the time interval $T := [1/4v_2, 3/4v_2]$, during which runner 2 is for the first time in the distant region $[\frac{1}{4}, \frac{3}{4}]$. For $r = 1, 3$, let $T_r = \{t \in [0, 1) : \langle tv_r \rangle \in [\frac{1}{4}, \frac{3}{4}]\}$.

If $T \setminus (T_1 \cup T_3) = \emptyset$, then use (1) with the defined prejump x_1 , an arbitrary $t_2 \in T \setminus (T_1 \cup T_3)$, and $\alpha = 1$: $\langle (t_1 + t_2)v \rangle = (\frac{1}{2}, 0, \frac{1}{2}) + \langle t_2 v \rangle$. Since 2 is the only distant runner at time t_2 , $\{1, 2, 3\}$ is distant at time $t_1 + t_2$.

We may now assume $T \subseteq T_1 \cup T_3$. Suppose that $T \subseteq T_i$ for some $i \in \{1, 3\}$. Then T is contained in one of the closed intervals comprising T_i , which implies $v_2 \geq v_i$. Furthermore, i first becomes distant no later than 2 does, so $v_2 \leq v_i$ which contradicts $v_2 \neq v_i$.

Thus $T \subseteq T_1 \cup T_3$, $T \cap T_i \neq \emptyset$ ($i = 1, 3$). Both $T \cap T_1$ and $T \cap T_3$ consist of disjoint closed intervals and their union is T . Hence, $\emptyset \neq (T \cap T_1) \cap (T \cap T_3) = T \cap T_1 \cap T_3$, and we are done. ■

4. THE CASE $k = 4$

Before completing the proof of Theorem 2.2, we set some notation and present two more prejump facts which hold true whenever $k+1$ is prime. The notation $a|b$ means that a divides b . For fixed $k \geq 2$ we partition the circle $C = [0, 1)$ as $\{0\} \cup C_1 \cup C_2$, where

$$C_1 := \left(0, \frac{1}{k+1}\right) \cup \left(\frac{k}{k+1}, 1\right) \cup \left\{ \frac{1}{k+1}, \frac{2}{k+1}, \dots, \frac{k}{k+1} \right\},$$

$$C_2 := \left(\frac{1}{k+1}, \frac{2}{k+1}\right) \cup \left(\frac{2}{k+1}, \frac{3}{k+1}\right) \cup \dots \cup \left(\frac{k-1}{k+1}, \frac{k}{k+1}\right).$$

Given a speed vector $v \in \mathbb{N}^k$ and a position $x \in X = X(v)$, we define $D := \{r \in R: (k+1) | v_r\}$ and partition the runners R as $R_0(x) \cup R_1(x) \cup R_2(x)$, where

$$\begin{aligned} R_0(x) &:= D \cup \{r \in R: x_r = 0\}, \\ R_1(x) &:= \{r \in R \setminus D: x_r \in C_1\}, \\ R_2(x) &:= \{r \in R \setminus D: x_r \in C_2\}. \end{aligned}$$

(3) Let $k+1$ be prime, and suppose there exists $x \in X$ in which D is distant and $|R_2(x)| < |R_0(x)|$. Then there exists a time when R is distant.

Proof. We consider the list of k positions $\langle x + j/(k+1)v \rangle$ ($1 = 1, 2, \dots, k$). Since $k+1$ is prime, we have

$$\begin{aligned} \left\langle \frac{1}{k+1} v_r \right\rangle &= \dots = \left\langle \frac{k}{k+1} v_r \right\rangle = 0 && \text{if } r \in D, \\ \left\{ \left\langle \frac{1}{k+1} v_r \right\rangle, \dots, \left\langle \frac{k}{k+1} v_r \right\rangle \right\} &= \left\{ \frac{1}{k+1}, \dots, \frac{k}{k+1} \right\} && \text{if } r \in R \setminus D. \end{aligned}$$

Using this, it is straightforward to check that, for $m=0, 1, 2$, each runner in $R_m(x)$ is distant in exactly $k-m$ of the listed positions. Thus, there are at most $|R_1(x)| + 2 |R_2(x)|$ positions in the list in which R is not distant. If $|R_2(x)| < |R_0(x)|$ then $|R_1(x)| + 2 |R_2(x)| < k$, so R is distant in at least one of the k listed positions. ■

Here is an easy corollary.

(4) Suppose that $k+1$ is prime and the only speed which it divides is v_2 . If there exists $d \in \mathbb{N}$ dividing at least $k/2$ different speeds, but not dividing v_2 , then there exists a time when R is distant.

Proof. Let $R' := \{r \in R: d | v_r\}$. Since $d \geq 2$ and $2 \notin R'$, there exists $j \in \{0, \dots, d-1\}$ such that runner 2 is distant in $x := \langle (j/d)v \rangle$. We have that $x_r = 0$ for each $r \in R'$, so $R_0(x) \supseteq \{2\} \cup R'$, and therefore $|R_0(x)| \geq 1 + |R'| > k/2 = |R|/2$, whence $|R_0(x)| > |R_2(x)|$. Since $D = \{2\}$ is distant, we are done by (3). ■

Proof of Theorem 2.2. We assume $k=4$, $R = \{1, 2, 3, 4\}$, all speeds are distinct and have no common prime factor. Consider the (proper) subset $D = \{r \in R: 5 | v_r\}$. If $|D| = 0$, then R is distant at time $\frac{1}{5}$. Suppose $2 \leq |D| \leq 3$. By induction on k there exists a position y where D is distant. Either we are done at y , or some runner in $R \setminus D$ is not distant, whence $|R_0(y)| + |R_1(y)| \geq |D| + 1 \geq 3$, so $|R_2(y)| \leq 1$, whereas $|R_0(y)| \geq |D| \geq 2 > 1 \geq |R_2(y)|$ and

we are done by (3). We henceforth assume $D = \{2\}$, whence $2 \in R_0(x)$ for every position x .

If no runner is faster than 2, then at time $1/5v_2$, 2 is the only distant runner, whence $|R_2(v/5v_2)| = 0$, $|R_0(v/5v_2)| = 1$, and we are again done by (3). We thus assume $v_1 > v_2, v_3, v_4$.

At least one of v_3, v_4 , say v_3 , is not equal to $v_1 - v_2$. Since v_2, v_3 are distinct and less than v_1 , the assumptions $v_3 \neq v_2$ and $v_3 \neq v_1 - v_2$ imply $v_3 \not\equiv \pm v_2 \pmod{v_1}$. If $d := \gcd(v_1, v_3) > 1$, then if d divides v_2 , we are done by (2); if it does not, we are done by (4).

Thus we can assume $\gcd(v_1, v_3) = 1$. Then there exists $\alpha \in \mathbb{N}$, $\alpha v_3 \equiv 1 \pmod{v_1}$. Let x be the position at time α/v_1 . We have $x_1 = 0$ and $x_3 = 1/v_1 < 1/v_2 \leq 1/5$, so $1, 2 \in R_0(x)$ and $3 \in R_1(x)$. If $D = \{2\}$ is distant in x , then we are done by (3) since $1, 2 \in R_0(x)$, whereas $3 \in R_1(x)$, so $|R_2(x)| \leq 1$. So we may assume 2 is not distant in x .

We notice two facts. First, the distance of x_2 from 0 is at least twice that of x_3 (this follows from $v_2 \not\equiv 0, \pm v_3 \pmod{v_1}$ and $\gcd(\alpha, v_1) = 1$, which implies $x_2 = \langle \alpha/v_1 \rangle \neq 0, \pm 1/v_1$, whence $x_2 \in [2/v_1, 1 - 2/v_1]$.) Second, if a runner has distance $\delta \leq 1/4$ from 0 in some position $z \in X$, then it has distance 2δ in position $\langle 2z \rangle$. Let x' be the first position in the sequence $\langle 2x \rangle, \langle 4x \rangle, \langle 8x \rangle, \dots$ in which 2 is distant. As before, $1, 2 \in R_0(x')$, whereas, by the two facts and the minimality in the choice of $x', x'_3 \in (0, 1/5)$ so $3 \in R_1(x')$, and we are again done by (3). ■

ACKNOWLEDGMENTS

The authors thank Andrew Odlyzko for directing them to the papers of Jüing Wills and Thomas Cusick, and they thank Martin Loebel for helpful discussions. The authors gratefully acknowledge the stimulating mathematical environments in Grenoble, Tel-Aviv, Vancouver, and Waterloo, where this work was done.

REFERENCES

1. U. Betke and J. M. Wills, Untere Schranken für zwei diophantische Approximations-Funktionen, *Monatsch. Math.* **76** (1972), 214–217.
2. Y. G. Chen, On a conjecture about diophantine approximations, I, *Acta Math. Sinica* **33** (1990), 712–717. [Chinese]
3. T. W. Cusick, View-obstruction problems, *Aequationes Math.* **9** (1973), 165–170.
4. T. W. Cusick, View-obstruction problems in n -dimensional geometry, *J. Combin. Theory Ser. A* **16** (1974), 1–11.
5. T. W. Cusick, View-obstruction problems, II, *Proc. Amer. Math. Soc.* **84** (1982), 25–28.
6. T. W. Cusick and C. Pomerance, View-obstruction problems, III, *J. Number Theory* **19** (1984), 131–139.
7. L. R. Ford and D. R. Fulkerson, Network flow and systems of representatives, *Canad. J. Math.* **10** (1958), 78–84.

8. A. J. Hoffman, Some recent applications of the theory of linear inequalities to external combinatorial analysis, in "Proc. Sympos. Appl. Math." (R. Bellman and M. Hall, Jr., Eds.), Vol. 10, pp. 113–127, Amer. Math. Soc., Providence, RI, 1960.
9. F. Jaeger, Nowhere-zero flow problems, in "Selected Topics in Graph Theory" (L. W. Beineke and R. Wilson, Eds.), Vol. 3, pp. 71–95, Academic Press, San Diego, CA, 1988.
10. F. Jaeger, Flows and generalized coloring theorems in graphs, *J. Combin. Theory Ser. B* **26** (1979), 205–216.
11. J. G. Oxley, "Matroid Theory," Oxford Univ. Press, Oxford, 1992.
12. P. D. Seymour, Decomposition of regular matroids, *J. Combin. Theory Ser. B* **28** (1980), 305–359.
13. P. D. Seymour, Somewhere-zero 6-flows, *J. Combin. Theory Ser. B* **30** (1981), 130–135.
14. P. D. Seymour, Nowhere-zero flows: Appendix: Colouring, stable sets and perfect graphs, in "Handbook of Combinatorics" (R. Graham, M. Grötschel, and L. Lovász, Eds.), Vol. 1, pp. 289–299, Elsevier, Amsterdam, 1995.
15. M. Tarsi, Nowhere-zero flow and circuit covering in regular matroids, *J. Combin. Theory Ser. B* **39** (1985), 346–352.
16. W. T. Tutte, A contribution to the theory of chromatic polynomials, *Canad. J. Math.* **6** (1954), 80–91.
17. J. M. Wills, Zwei Sätze über inhomogene diophantische Approximation von Irrationalzahlen, *Monatsch. Math.* **71** (1967), 263–269.
18. J. M. Wills, Zur simultanen homogenen diophantischen Approximation, I, *Monatsch. Math.* **72** (1968), 254–263.
19. J. M. Wills, Zur simultanen homogenen diophantischen Approximation, II, *Monatsch. Math.* **72** (1968), 268–281.
20. J. M. Wills, Zur simultanen homogenen diophantischen Approximation, III, *Monatsch. Math.* **74** (1970), 166–171.
21. J. M. Wills, Zur simultanen homogenen diophantischen Approximation, *Zahlentheorie (Tagung, Math. Forschungsinst. Oberwolfach, 1970)*, Ber. Math. Forschungsinst., Oberwolfach, (No. 5) pp. 223–227, Bibliographisches Inst., Mannheim, 1971.
22. "Weissman Institute Public Relations, Booklet, 1994."