

Edge-connectivity augmentations of graphs and hypergraphs*

Zoltán Szigeti

Laboratoire G-SCOP, INPGrenoble, UJF, CNRS, 46 avenue Félix Viallet 38031 Grenoble Cedex, France

Summary. A. Frank (Augmenting graphs to meet edge-connectivity requirements, SIAM J. on Disc. Math. Vol. 5, No. 1 (1992) 22-53) developed a method to solve edge-connectivity augmentation problems. His paper has stimulated further research in a number of directions, including many interesting generalizations.

This paper surveys the current State of the Art on the edge-connectivity augmentation problem. Recent extensions of the problem are presented for undirected graphs, hypergraphs and more generally for set functions. Shortened proofs are provided for some of the results. A list of open problems is also presented.

1.1 Introduction

In this paper all graphs and hypergraphs are undirected, directed versions of the problems will not be treated here. By graphs and hypergraphs we mean multi-graphs and multi-hypergraphs, that is, parallel edges and parallel hyperedges are allowed, however loops are forbidden. All the problems here will concern edge-connectivity, that is, vertex-connectivity will not be considered. We have to emphasize immediately that in our problems we may add edges between adjacent vertices, we may even add parallel edges. We remark that without these assumptions, that is, when the starting and also the resulting graph must be simple, the problem is NP-complete [22]. The optimization problems will always be unweighted, the weighted version of the simplest problem already being NP-complete [12].

The basic problem: The starting point for introducing edge-connectivity problems is the problem of increasing the reliability of a telephone network. We may associate a graph to the network: the telephone centers and the connections between them are the vertices and the edges of the graph and the

* Some part of this work was done while the author was visiting the Research Institute for Discrete Mathematics, University of Bonn, Lennéstrasse 2, 53113. Bonn, Germany by an Alexander von Humboldt fellowship.

reliability of the network corresponds to the edge-connectivity of this graph. As a natural requirement, we may wish to increase the reliability of the network by constructing new connections between the centers. The optimization problem in the language of graphs is the global edge-connectivity augmentation problem in graphs, namely: Given a graph $G = (V, E)$ and a positive integer k , what is the minimum number γ of new edges whose addition results in a k -edge-connected graph? We show how to solve this problem, in doing so we introduce the key ideas to be applied throughout this paper.

The lower bound: First we provide a lower bound on γ . Suppose that G is not k -edge-connected. This is because there is a set X of degree $d(X)$ less than k . Then the deficiency of X is $k - d(X)$, that is, we must add at least $k - d(X)$ edges between X and $V - X$. Let $\{X_1, \dots, X_l\}$ be a subpartition of V . The deficiency of this subpartition is the sum of the deficiencies of the X_i 's. By adding a new edge we may decrease the deficiency of at most two X_i 's so we may decrease the deficiency of the subpartition by at most two, hence we obtain the following lower bound:

$$\gamma \geq \alpha := \lceil \text{half of the maximum deficiency of a subpartition of } V \rceil. \quad (1.1)$$

The minimax theorem (see Theorem 3), due to Watanabe and Nakamura [36], saying that this lower bound can always be achieved, can be proved as follows:

Frank's algorithm: (1) *Minimal extension:* First, add a new vertex s to G and connect it to each vertex of G by k edges. The resulting graph is k -edge-connected in V . Secondly, delete as many new edges as possible preserving k -edge-connectivity in V . The graph obtained is denoted by $G' = (V, F')$. If the degree of s in G' is odd, then add an arbitrary new edge incident to s . Then we have a graph $G'' = (V + s, E \cup F'')$ that is k -edge-connected in V so that the degree of s is even.

(2) *Splitting off:* Now we will use the main operation of this paper, called splitting off. Splitting off a pair of edges sr, st incident to s means that we delete these two edges and we add a new edge rt . Applying Lovász's theorem 2(a), split off edge pairs incident to s , preserving k -edge-connectivity in V , as far as the degree of s becomes zero. This way we obtain a k -edge-connected graph $G^* = (V, E \cup F)$ with $|F| = \frac{|F''|}{2} = \lceil \frac{|F'|}{2} \rceil$.

Optimality: The optimality will be proved by the existence of a subpartition of V whose deficiency provides equality in (1.1). In G' , no edge incident to s can be deleted without violating k -edge-connectivity in V , so each edge $e \in F'$ enters a maximal proper subset X_e in V of degree k , that is, $d_G(X_e) + d_{F'}(X_e) = k$. By the submodularity of the degree function d , these sets provide a subpartition $\{X_1, \dots, X_l\}$ of V , for which we do have equality in (1.1):

$$\gamma \leq |F| = \lceil \frac{|F'|}{2} \rceil = \lceil \frac{1}{2} \sum_1^l d_{F'}(X_i) \rceil = \lceil \frac{1}{2} \sum_1^l (k - d_G(X_i)) \rceil \leq \alpha \leq \gamma.$$

Minimal extension: The above method of Frank has two main phases: the first one - minimal extension - consists of the first two steps, while the second one is the splitting off.

The first phase is in fact the construction of a graph $H = (V + s, F)$ with a minimum number of edges such that each edge of H is incident to s and H covers the deficiency function of G , that is, for every vertex set $X \subset V$, the number of edges of H leaving X is at least the deficiency of X in G with respect to k .

Frank proved in [15] that this can be done not only for the deficiency function of a graph: such an optimal graph H can be constructed that covers a symmetric skew-supermodular function. This result (Theorem 1 in this paper) is not explicitly presented in [15], it was published in [1].

This general result on extension implies that for an edge-connectivity augmentation problem, if the corresponding splitting off exists, then the optimization problem can be solved. Thus we will concentrate on splitting off results in this paper.

Generalizations: The above mentioned basic problem (that is the problem of augmenting global edge-connectivity of a graph by adding graph edges) and its solution capture already the most important ingredients of the theory and they provide a point of departure for studying more complex edge-connectivity augmentation problems. Lots of generalizations of Watanabe and Nakamura's result will be presented here. This paper is divided in three parts: results on graphs, on hypergraphs and on set functions. The results in the different parts are intimately related, we will see that a great number of results on graphs can be generalized for hypergraphs, which in turn can sometimes be further extended to "connectivity" functions.

The first part contains the following generalizations in *graphs*:

- local edge-connectivity augmentation [15],
- global edge-connectivity augmentation over symmetric parity families [32],
- node to area global edge-connectivity augmentation [19],
- global edge-connectivity augmentation by attaching stars [14],
- local edge-connectivity augmentation by attaching stars [24],
- global edge-connectivity augmentation with partition constraint [2].

We present in the second part generalizations in *hypergraphs*:

- global edge-connectivity augmentation in hypergraphs by adding graph edges [3],
- global edge-connectivity augmentation in hypergraphs by adding uniform hyperedges [26],
- local edge-connectivity augmentation in hypergraphs by adding graph edges (we mention at once that this problem is NP-complete [11]),
- local edge-connectivity augmentation in hypergraphs by adding a hypergraph of minimum total size [31].

The deficiency function with respect to global (resp. local) edge-connectivity in graphs and in hypergraphs is symmetric and crossing supermodular (resp. skew-supermodular). The third part is devoted to generalizations on such *set functions*:

- covering a symmetric crossing supermodular set function by a graph [6],
- covering a symmetric crossing supermodular set function by a uniform hypergraph [26],
- covering a symmetric skew-supermodular set function by a graph (this problem is NP-complete [27]),
- covering a symmetric semi-monotone set function by a graph [20],
- covering a symmetric skew-supermodular set function by a hypergraph of minimum total size [31].

The main contribution of this paper is to call the reader's attention to Theorem 1 of Frank, to survey the results on the edge-connectivity augmentation problem, to provide short proofs for Theorem 12 on detachments satisfying local edge-connectivity requirements, and for Theorem 14 on partition constrained splitting off preserving global edge-connectivity, and finally, to present some open problems of the theory.

We finish this introduction by emphasizing that we do not attempt to cover all topics of the field, e.g. we do not focus on the design of efficient algorithms, but instead, we concentrate on minimax results of the area.

For further topics, such as Local edge-connectivity augmentation of mixed graphs, Global edge-connectivity augmentation preserving simplicity, Global edge-connectivity augmentation in a graph by adding edges within the members of a partition, Successive edge-connectivity augmentation, Simultaneous global edge-connectivity augmentation, we refer to [1], [4], [5], [9], [23].

1.2 Definitions

This section is divided in three parts: definitions on graphs, on hypergraphs and finally on set functions.

Graph: Let $G = (V, E)$ be a graph. Recall that parallel edges are allowed. The set of all subpartitions of V will be denoted by $\mathcal{S}(V)$. For a vertex $v \in V$, $\Gamma(v)$ denotes the neighbours of v . For $X, Y \subset V$, G/X denotes the graph obtained from G by contracting X into one vertex, while the resulting parallel edges are kept, $d(X, Y)$ (resp. $\bar{d}(X, Y)$) denotes the number of edges between $X - Y$ and $Y - X$ (resp. $X \cap Y$ and $V - (X \cup Y)$), $d(X) = d(X, V - X)$. The set of edges leaving X is called a **cut** and is denoted by $\delta(X)$, that is, $d(X) = |\delta(X)|$. It is well-known and easy to check that, for all $X, Y \subseteq V$, (1.2) and (1.3) are satisfied.

$$d(X) + d(Y) = d(X \cap Y) + d(X \cup Y) + 2d(X, Y), \quad (1.2)$$

$$d(X) + d(Y) = d(X - Y) + d(Y - X) + 2\bar{d}(X, Y). \quad (1.3)$$

The **local edge-connectivity** between two different vertices x and y of G is defined by $\lambda_G(x, y) = \min\{d_G(X) : x \in X, y \notin X\}$, while $\lambda_G(x, x) = +\infty$. By Menger's theorem, $\lambda_G(x, y)$ is the maximum number of edge disjoint paths in G between x and y . Let $G = (U, E)$ be a graph. For $X \subset U$, $x, y \in U - X$, $s \in U$, $u, v \in U - s$ we have

$$\lambda_{G/X}(x, y) \geq \lambda_G(x, y), \quad (1.4)$$

$$\lambda_{G-s}(u, v) \geq \lambda_G(u, v) - \lfloor \frac{d_G(s)}{2} \rfloor. \quad (1.5)$$

Indeed, if a cut Q separates x and y in G/X then Q also separates them in G and (1.4) follows. On the other hand, if \mathcal{P} is a set of $\lambda_G(u, v)$ edge-disjoint paths in G then at most $\lfloor \frac{d_G(s)}{2} \rfloor$ of them may contain the vertex s and hence the other paths of \mathcal{P} belong to $G - s$ and (1.5) follows.

A graph $G = (V, E)$ is called **k -edge-connected in U** (for some $k \in \mathbb{Z}_+$ and $U \subseteq V$) if $\lambda_G(x, y) \geq k \forall x, y \in U$. This definition will be usually used for $U = V$ or $V - s$ with a specified vertex s of V , in which case it is equivalent to (1.6). $\lambda(G)$ denotes the global edge-connectivity of G that is the maximum integer k so that G is k -edge-connected. Given a symmetric function $r : V \times V \rightarrow \mathbb{Z}_+$, we say that G is **r -edge-connected** if (1.7) is satisfied.

$$d_G(X) \geq k \quad \forall \emptyset \neq X \subset U, \quad (1.6)$$

$$\lambda_G(u, v) \geq r(u, v) \quad \forall u, v \in V. \quad (1.7)$$

Let $G = (V + s, E)$ be a graph. By **splitting off** two edges sr, st we mean the operation that replaces sr, st by a new edge rt , the new graph will be denoted by G_{rt} . Note that we do not allow loops so if $r = t$, then the resulting loop must be deleted. A pair sr, st is called **k -admissible** if G_{rt} is k -edge-connected in V and it is called **λ -admissible** if (1.8) is satisfied. On several occasions, the splitting off will be called k -admissible instead of the edge pair. A sequence of splittings off is said to be **complete** if the degree of s becomes 0. A complete splitting off is called **k -admissible (λ -admissible)** if each splitting off in its sequence is k -admissible (λ -admissible, respectively).

$$\lambda_{G_{rt}}(x, y) \geq \lambda_G(x, y) \quad \forall x, y \in V. \quad (1.8)$$

Hypergraph: Let $\mathcal{G} := (V, E)$ be a **hypergraph**, that is E is a multiset of subsets of V . The element of E are called **hyperedges**. Note that parallel hyperedges are allowed. The above definitions of the degree function, local edge-connectivity and k -edge-connectivity can be naturally generalized for hypergraphs. Indeed, let $\mathcal{G} := (V, E)$ be a hypergraph. The degree $d_{\mathcal{G}}(X)$

of a vertex set X is defined as the number of hyperedges intersecting X and $V - X$. The **local edge-connectivity** between two different vertices x and y of \mathcal{G} is defined by $\lambda_{\mathcal{G}}(x, y) = \min\{d_{\mathcal{G}}(X) : x \in X, y \notin X\}$, while $\lambda_{\mathcal{G}}(x, x) = +\infty$. \mathcal{G} is called **k -edge-connected in U** (for some $k \in \mathbb{Z}_+$ and $U \subseteq V$) if $\lambda_{\mathcal{G}}(x, y) \geq k \forall x, y \in U$. We say that a hypergraph \mathcal{H} **covers** a function p if

$$d_{\mathcal{H}}(X) \geq p(X) \quad \forall \emptyset \neq X \subset V. \quad (1.9)$$

Call a set X of V **tight** if $d_{\mathcal{H}}(X) = p(X)$. $\mathbf{c}(\mathcal{G})$ denotes the number of connected components of \mathcal{G} . A hypergraph is **r -uniform** if each hyperedge is of size r . A graph is a 2-uniform hypergraph. \mathcal{G} is called a **2-3 hypergraph** if each hyperedge is of size two or three. The operation $\Delta - Y$ replaces a given 3-hyperedge abc by the star qa, qb, qc of a new vertex q of degree three. The operation $Y - \Delta$ replaces the star qa, qb, qc of a given vertex q of degree three by a new 3-hyperedge abc . It is easy to check that the local edge-connectivities between the original vertices do not change after a $\Delta - Y$ or a $Y - \Delta$ operation.

Set function: Let $p : 2^V \rightarrow \mathbb{Z} \cup \{-\infty\}$ be a set function. The function p is called **supermodular (crossing supermodular)** if (1.10) holds for all $X, Y \subseteq V$ (for all $X, Y \subseteq V$ that are **crossing** that is $X \cap Y, X - Y, Y - X, V - (X \cup Y) \neq \emptyset$) and p is called **skew-supermodular** if at least one of (1.10) and (1.11) hold for all $X, Y \subseteq V$. We say that p is **symmetric** if (1.12) is satisfied for each $X \subseteq V$. A function $b : 2^V \rightarrow \mathbb{Z}$ is called **submodular** if $-b$ is supermodular. Note that the degree function $d_G(X)$ of a graph G is symmetric and, by (1.2), it is submodular.

$$p(X) + p(Y) \leq p(X \cap Y) + p(X \cup Y), \quad (1.10)$$

$$p(X) + p(Y) \leq p(X - Y) + p(Y - X), \quad (1.11)$$

$$p(X) = p(V - X). \quad (1.12)$$

Given a symmetric function $r : V \times V \rightarrow \mathbb{Z}_+$, let us define $\mathbf{R}(X) := \max\{r(x, y) : x \in X, y \in V - X\}$. It is known that R is skew-supermodular (see e.g. in [16]).

Symmetric crossing supermodular functions generalize the deficiency function of a graph G concerning global edge-connectivity (that is the function $k - d_G(X)$ is symmetric crossing supermodular) and symmetric skew-supermodular functions generalize the deficiency function of a graph G concerning local edge-connectivity (that is the function $R(X) - d_G(X)$ is symmetric skew-supermodular).

1.3 Minimal extension

We start this section with a typical lemma that shows how to use the skew-supermodularity of a function. Recall that if a graph H covers a set function p , then a set X of V is called tight if $d_H(X) = p(X)$.

Lemma 1. *Let $p : 2^V \rightarrow \mathbb{Z} \cup \{-\infty\}$ be a symmetric skew-supermodular set function. Let $H = (V + s, E)$ be a graph that covers p . If X and Y are tight sets and $X \cap Y \neq \emptyset$, then either (a) $X \cap Y$ and $X \cup Y$ are tight, or (b) $X - Y$ and $Y - X$ are tight and $\bar{d}(X, Y) = 0$.*

Proof. We may suppose that $X - Y \neq \emptyset \neq Y - X$ because otherwise (a) is trivially satisfied.

If p satisfies (1.11) for X and Y then, by (1.3) and (1.9),

$$\begin{aligned} p(X) + p(Y) &= d_H(X) + d_H(Y) \\ &= d_H(X - Y) + d_H(Y - X) + 2\bar{d}(X, Y) \\ &\geq p(X - Y) + p(Y - X) + 0 \\ &\geq p(X) + p(Y), \end{aligned}$$

so we have equality everywhere, implying (b).

Otherwise, $X \cup Y \neq V$ and p satisfies (1.10) for X and Y . Then, by (1.2) and (1.9),

$$\begin{aligned} p(X) + p(Y) &= d_H(X) + d_H(Y) \\ &\geq d_H(X \cap Y) + d_H(X \cup Y) \\ &\geq p(X \cap Y) + p(X \cup Y) \\ &\geq p(X) + p(Y), \end{aligned}$$

so we have equality everywhere, implying (a). \square

All the augmentation results of this paper will be obtained by applying the following general result of Frank and some suitable splitting off theorem. For the sake of completeness we provide the proof of this theorem.

Theorem 1 (Frank [15], [1]). *Let $p : 2^V \rightarrow \mathbb{Z} \cup \{-\infty\}$ be a symmetric skew-supermodular function. Then the edgeless graph on V can be extended to a graph H by adding a new vertex s and γ edges incident to s so that H covers p if and only if*

$$\sum_{X \in \mathcal{X}} p(X) \leq \gamma \quad \forall \mathcal{X} \in \mathcal{S}(V). \tag{1.13}$$

Proof. Suppose that $H = (V + s, E)$ covers p , each edge of H is incident to s and $d_H(s) \leq \gamma$. Then, for any subpartition \mathcal{X} of V , (1.13) is satisfied by

$$\sum_{X \in \mathcal{X}} p(X) \leq \sum_{X \in \mathcal{X}} d_H(X) \leq d_H(s) \leq \gamma.$$

Now suppose that (1.13) is satisfied. The desired graph is constructed as follows. First add a new vertex s to V and connect it to each vertex of V by $\max\{p(X) : X \subset V\}$ edges. Then, of course, this graph covers p . Secondly,

delete as many edges as possible preserving that p is covered. Let H be the graph obtained. It remains to show that $d_H(s) \leq \gamma$.

No edge of H can be deleted, that is, each edge of H enters a tight set. Thus there exists a set $\mathcal{X} := \{X_1, \dots, X_l\}$ of tight sets so that each edge enters some set X_i and $\sum_1^l |X_i|$ is minimal.

We claim that \mathcal{X} is a subpartition of V . Suppose that $X_i \cap X_j \neq \emptyset$ for some $X_i, X_j \in \mathcal{X}$. By Lemma 1, either $X_i \cup X_j$ is tight, hence X_i and X_j can be replaced by $X_i \cup X_j$ or $X_i - X_j$ and $X_j - X_i$ are tight and $\bar{d}(X_i, X_j) = 0$ (that is no edge enters $X_i \cap X_j$), so X_i and X_j can be replaced by $X_i - X_j$ and $X_j - X_i$. In both cases we obtained a contradiction to the fact that $\sum_1^l |X_i|$ is minimal.

Thus, $\mathcal{X} \in \mathcal{S}(V)$, so we are done because, by (1.13),

$$d_H(s) = \sum_{X \in \mathcal{X}} d_H(X) = \sum_{X \in \mathcal{X}} p(X) \leq \gamma. \quad \square$$

1.4 Graphs

In this section we will present the problems on edge-connectivity augmentation in graphs and their solutions. As we mentioned in the introduction, the basic tool is splitting off. Each subsection is devoted to one problem, and it is divided in two parts : results on splitting off and then, applying Theorem 1, we read out the minimax result on augmentation.

1.4.1 Global edge-connectivity I

This section is about k -edge-connectivity for some $k \geq 2$. First we consider the operation splitting off: here the graph $G = (V + s, E)$ is k -edge-connected in V and we wish to reduce the graph (by splitting off an edge pair incident to s) in such a way that the graph remains k -edge-connected in V . Then we consider the augmentation problem where the aim is to make a graph k -edge-connected by adding new edges.

Splitting off preserving global edge-connectivity

The first result on splitting off is due to Lovász [28]. It concerns global edge-connectivity, namely it provides a sufficient condition for the existence of a k -admissible splitting off. More precisely, Lovász showed that if $k \geq 2$ and the degree of the special vertex s is even, then each edge belongs to a k -admissible pair at s . In [2] it was shown that each edge belongs to many k -admissible pairs.

Theorem 2. *Let $G = (V + s, E)$ be a k -edge-connected graph in V with $k \geq 2$ and $d(s)$ is even. Then :*

- (a) (**Lovász [28]**) *each edge st belongs to a k-admissible pair at s.*
- (b) (**Bang-Jensen, Gabow, Jordán, Szigeti [2]**) *each edge st belongs to at least $\frac{d(s)}{2}$ (resp. $\frac{d(s)}{2} - 1$) k-admissibles pairs at s if k is even (resp. odd).*

B. Fleiner [14], and independently Bang-Jensen and Jackson [3] proved that Theorem 2(a) is true for 2-3 hypergraphs containing no 3-hyperedges incident to s . The special case of Theorem 2(b), when G is Eulerian, was proved earlier by Jackson [21].

We must emphasize that the above theorem is true only if $k \geq 2$. Let us consider the following example for $k = 1$: let the graph $G = (V + s, E)$ be the star of the vertex s of degree four. Then G is connected in V and there is no complete 1-admissible splitting off. To avoid this problem we will usually suppose that the connectivity requirement is at least two.

Augmentation of global edge-connectivity by adding edges

The problem of **global edge-connectivity augmentation in graphs**, already introduced and also solved in the introduction, is the following: *Given a graph $G = (V, E)$ and an integer k , what is the minimum number γ of new edges whose addition results in a k-edge-connected graph?* In other words, we are looking for

$$\begin{aligned} \gamma &:= \min\{|E'| : d_{G+E'}(X) \geq k \ \forall \emptyset \neq X \subset V\} \\ &= \min\{|E'| : d_{(V,E')}(X) \geq k - d_G(X) \ \forall \emptyset \neq X \subset V\}. \end{aligned}$$

As the function $p(X) = k - d_G(X)$ is symmetric and, by (1.2), skew-supermodular, Theorems 1 and 2(a) imply the following theorem. (Cai and Sun [8] also proved this result later using some splitting off technique.)

Theorem 3 (Watanabe and Nakamura [36]). *Let $G = (V, E)$ be a graph and $k \geq 2$. Then G can be made k-edge-connected by adding at most γ new edges if and only if*

$$\sum_{X \in \mathcal{X}} (k - d_G(X)) \leq 2\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V). \tag{1.14}$$

We point out again that the case when $k = 1$ does not fit into this framework. It is obvious that in this case we have to add $l - 1$ new edges, where l is the number of connected components of the graph.

1.4.2 Local edge-connectivity I

In the above section we were interested in k -edge-connectivity, that is, in the minimum value of the local edge-connectivities over all pairs of vertices. This section concerns the problem where each value counts not just the minimum, that is, we wish to add new edges to a graph so that the local edge-connectivity be greater than or equal to a given requirement for each pair of vertices.

Splitting off preserving local edge-connectivity

In this section we summarize results on splitting off preserving local edge-connectivity. Mader [29] generalized Lovász's result Theorem 2(a) for local edge-connectivity by showing that a λ -admissible pair always exists if the degree of the vertex s is different from 3 and roughly speaking G is 2-edge-connected. This result implies that at most three edges incident to s belong to no λ -admissible pair. Frank [16] improved this by showing that at most one edge incident to s belongs to no λ -admissible pair. In [33] we characterized this edge. Since not every edge belongs to a λ -admissible pair, the best we may hope for is that there exists at least one edge that belongs to many λ -admissible pairs. In [33] it is shown that the correct number is $\lfloor \frac{d(s)}{3} \rfloor$ and examples show that this result is best possible.

Theorem 4. *Let $G = (V + s, E)$ be a connected graph so that $d(s) \neq 3$ and no cut edge is incident to s .*

- (a) (Mader [29]) *There exists a λ -admissible pair at s .*
- (b) (Frank [16]) *There exist $\lfloor \frac{d(s)}{2} \rfloor$ disjoint λ -admissible pairs at s . (Hence at most one edge incident to s belongs to no λ -admissible pair.)*
- (c) (Szigeti [33]) *An edge st belongs to no λ -admissible pair if and only if $d(s)$ is odd and there exist two disjoint sets $C_1, C_2 \subset V - t$ such that $d(C_i) = R(C_i)$ and $d(s, C_i) = \frac{d(s)-1}{2}$ for $i = 1, 2$. Moreover, for every $c_1 \in C_1 \cap \Gamma(s), c_2 \in C_2 \cap \Gamma(s)$, $\{sc_1, sc_2\}$ is a λ -admissible pair.*
- (d) (Szigeti [33]) *There exists an edge belonging to at least $\lfloor \frac{d(s)}{3} \rfloor$ λ -admissible pairs at s .*

Augmentation of local edge-connectivity by adding edges

The problem of **local edge-connectivity augmentation in graphs** is defined as follows: *Given a graph G and a symmetric requirement function $r : V \times V \rightarrow \mathbb{Z}_+$, what is the minimum number γ of new edges whose addition results in an \mathbf{r} -edge-connected graph?*

Note that, by taking r to be equal to k for each pair of vertices, this problem contains, as a special case, the global edge-connectivity augmentation problem in graphs.

Recall that $R(X) = \max\{r(x, y) : x \in X, y \in V - X\}$ is symmetric and skew-supermodular. We can reformulate the problem as follows: we look for

$$\begin{aligned} \gamma &:= \min\{|E'| : \lambda_{G+E'}(u, v) \geq r(u, v) \forall u, v \in V\} \\ &= \min\{|E'| : d_{G+E'}(X) \geq R(X) \forall \emptyset \neq X \subset V\} \\ &= \min\{|E'| : d_{(V, E')}(X) \geq R(X) - d_G(X) \forall \emptyset \neq X \subset V\}. \end{aligned}$$

As $p(X) = R(X) - d_G(X)$ is a symmetric, skew-supermodular function, Theorems 1 and 4 imply the following theorem.

Theorem 5 (Frank [15]). *Let $G = (V, E)$ be a graph and $2 \leq r(u, v) \in \mathbb{Z} \forall u, v \in V$. Then G can be made \mathbf{r} -edge-connected by adding at most γ new edges if and only if*

$$\sum_{X \in \mathcal{X}} (R(X) - d_G(X)) \leq 2\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V). \quad (1.15)$$

Note that the special case of Theorem 5 when $r(u, v) = k \forall u, v \in V$ is exactly Theorem 3.

1.4.3 Symmetric parity families

A family \mathcal{F} of subsets of V is called a **symmetric parity family** if it satisfies the following three properties. (i) $\emptyset, V \notin \mathcal{F}$, (ii) if $A \in \mathcal{F}$, then $V - A \in \mathcal{F}$, (iii) if $A, B \notin \mathcal{F}$ and $A \cap B = \emptyset$, then $A \cup B \notin \mathcal{F}$. Let $T \subseteq V$ be a set of even cardinality. A set X is called **T-odd** if $|T \cap X|$ is odd. If $G = (V, E)$ is a connected graph then (G, T) is called a **graft**. A cut $\delta_G(X)$ is called a **T-cut** if X is T-odd, more generally $\delta_G(X)$ is called an **\mathcal{F} -cut** if $X \in \mathcal{F}$. The most important examples of parity families are $\mathcal{F} := 2^V - \{\emptyset, V\}$ and $\mathcal{F} := \{X \subseteq V : X \text{ is T-odd}\}$.

In this section we will deal with \mathcal{F} -cuts.

Splitting off preserving global edge-connectivity over symmetric parity families

Theorem 4(a) implies easily the following.

Theorem 6 (Szigeti [32]). *Let $G = (V + s, E)$ be a graph so that $d(s) > 0$ is even and let \mathcal{F} be a symmetric parity family on V . Suppose that for some $k \geq 2$,*

$$d(X) \geq k \quad \forall X \in \mathcal{F}. \quad (1.16)$$

Then there exists a pair of edges incident to s that can be split off without violating (1.16).

Augmentation of global edge-connectivity over symmetric parity families

In this section we solve the following **global edge-connectivity augmentation problem over a symmetric parity family**: *Given a graph $G = (V, E)$, a symmetric parity family \mathcal{F} on V and an integer k , what is the minimum number γ of edges whose addition results in a graph in which each \mathcal{F} -cut contains at least k edges?*

A special case is the minimum T-cut augmentation problem: how many new edges must be added to a graph so that the minimum T-cut contains at

least k edges? It also contains, as a special case, the global edge-connectivity augmentation problem in graphs.

For a symmetric parity family \mathcal{F} , a graph $G = (V, E)$ and $k \in \mathbb{Z}^+$, let $p(X) := k - d_G(X)$ if $X \in \mathcal{F}$, and $-\infty$ otherwise. The problem of edge-connectivity augmentation over a symmetric parity family can be reformulated as follows: what is the value

$$\begin{aligned} \gamma &:= \min\{|E'| : d_{G+E'}(X) \geq k \ \forall X \in \mathcal{F}\} \\ &= \min\{|E'| : d_{(V, E')}(X) \geq p(X) \ \forall X \subseteq V\} \end{aligned}$$

It is easy to see that p is a symmetric skew-supermodular function, so Theorems 1 and 6 provide at once the following theorem.

Theorem 7 (Szigeti [32]). *For a graph $G = (V, E)$, a symmetric parity family \mathcal{F} on V and an integer $k \geq 2$, the minimum cardinality of an \mathcal{F} -cut can be augmented to k by adding at most γ edges if and only if*

$$\sum_{i=1}^l (k - d(X_i)) \leq 2\gamma \quad \forall \{X_1, \dots, X_l\} \in \mathcal{S}(V) \text{ with } X_i \in \mathcal{F}. \quad (1.17)$$

By applying Theorem 7, for $\mathcal{F} = 2^V - \{\emptyset, V\}$ we get Theorem 3, and, for \mathcal{F} being the set of T-odd subsets of V , we get the following theorem on T-cuts.

Theorem 8 (Szigeti [32]). *For any graft (G, T) , the minimum cardinality of a T-cut can be augmented to $k \geq 2$ by adding at most γ edges if and only if $\sum_{X \in \mathcal{X}} (k - d(X)) \leq 2\gamma$ for each subpartition \mathcal{X} of V into T-odd sets.*

1.4.4 Node to area edge-connectivity

The **node to area global edge-connectivity augmentation** problem can be defined as follows: *Given a graph $G = (V, E)$, a family \mathcal{A} of sets $A \subseteq V$ (called areas), and a requirement function $r : \mathcal{A} \rightarrow \mathbb{Z}_+$, add a minimum number $\mathbf{Opt}(r, G)$ of new edges to G so that the resulting graph contains $r(A)$ edge-disjoint paths from any area A to any vertex $v \notin A$.*

Note that, by taking just one vertex as the family of areas and k as the requirement for this vertex, then we get as a special case the global edge-connectivity augmentation problem in graphs.

Let us define $P_{\mathcal{A}}(X) = \max\{r(A) : A \in \mathcal{A}, A \cap X = \emptyset \text{ or } A \subseteq X\}$ if $V \neq X \neq \emptyset$ and $P_{\mathcal{A}}(V) = P_{\mathcal{A}}(\emptyset) = 0$ and $Q_{\mathcal{A}}(G) := \max\{\sum_{X \in \mathcal{X}} q_{\mathcal{A}}(X) : \mathcal{X} \in \mathcal{S}(V)\}$, where $q_{\mathcal{A}}(X) = P_{\mathcal{A}}(X) - d_G(X)$.

Now we can provide a lower bound for the optimal value, namely $\mathbf{Opt}(r, G) \geq \lceil \frac{Q_{\mathcal{A}}(G)}{2} \rceil$. The question is whether we have always equality here or not. Usually equality will hold, unless the graph contains a special configuration.

The node to area global edge-connectivity augmentation problem is a special case of the symmetric semi-monotone function covering problem treated in Section 1.6.2, so Theorem 29 implies the following. We mention that the above defined $P_{\mathcal{A}}(X)$ is the symmetric semi-monotone function to be covered.

Theorem 9 (Ishii, Hagiwara [19]). *Let $G = (V, E)$ be a graph, \mathcal{A} a family of sets $A \subseteq V$, and $r : \mathcal{A} \rightarrow \mathbb{Z}_+$ a requirement function so that $r(A) \neq 1 \forall A \in \mathcal{A}$. If G contains no \mathcal{A} -configuration, then $Opt(r, G) = \lceil \frac{Q_{\mathcal{A}}(G)}{2} \rceil$, otherwise $Opt(r, G) = \lceil \frac{Q_{\mathcal{A}}(G)}{2} \rceil + 1$.*

The definition of a \mathcal{A} -configuration can be found in [19], where it is called P-property. We mention that if $\mathcal{A} = \{v\}$ and $r(v) = k$, then no \mathcal{A} -configuration can exist, so Theorem 9 implies Theorem 3. We remark that without the condition $r(A) \neq 1 \forall A \in \mathcal{A}$, the problem is NP-complete, see Section 1.6.2.

1.4.5 Global edge-connectivity II

Let us return to global edge-connectivity. First we generalize the operation splitting off, by introducing detachment, and then we consider the problem where we wish again to make a graph k -edge-connected, but this time by attaching stars. The essential tool to solve this problem is exactly the operation detachment.

Detachments preserving global edge-connectivity

In this section we generalize the operation splitting off.

Let $G = (V + s, E)$ be a graph. A **degree specification** for s is a sequence $\mathbf{f}(s) = (d_1, \dots, d_p)$ of positive integers with $\sum_{j=1}^p d_j = d_G(s)$. An $\mathbf{f}(s)$ -**detachment** of G at s is the graph G' obtained from G by replacing s by a set s_1, \dots, s_p of independent vertices and distributing the edges incident to s among them in such a way that $d_{G'}(s_i) = d_i$ ($1 \leq i \leq p$). Note that all the other ends of the edges in G remain the same.

Let us mention that a splitting off can really be considered as a special case of detachment, namely if after a splitting off we subdivide the new edge then the new graph is a $(2, d(s) - 2)$ -detachment and this operation does not change the local edge-connectivities between the original vertices.

The following beautiful theorem of B. Fleiner characterizes graphs that have a k -edge-connected $\mathbf{f}(s)$ -detachment.

Theorem 10 (B.Fleiner [14]). *Let $G = (V + s, E)$ be a graph, $2 \leq k \in \mathbb{Z}$ and $\mathbf{f}(s) = (d_1, \dots, d_p)$ a degree specification for s with $d_i \geq 2 \forall i$. Then there exists an $\mathbf{f}(s)$ -detachment of G that is k -edge-connected in V if and only if*

$$G \text{ is } k\text{-edge-connected in } V, \tag{1.18}$$

$$G - s \text{ is } (k - \sum_1^p \lfloor \frac{d_i}{2} \rfloor)\text{-edge-connected.} \tag{1.19}$$

We note that the special case of Theorem 10, when each d_i is even, is equivalent to Theorem 2(a). Indeed, in this case the condition (1.19) is automatically satisfied by (1.5) and hence Theorem 10 implies Theorem 2(a). On the other hand, by Theorem 2(a), there exists a complete k -admissible splitting off, subdividing each new edge by a vertex and combining the suitable number of new vertices, an $f(s)$ -detachment is obtained which is k -edge-connected in V by (1.4), and hence Theorem 2(a) implies this special case of Theorem 10.

The proof technique of B. Fleiner [14] needed the more general framework of 2-3 hypergraphs. He proved that Theorem 10 is true for 2-3 hypergraphs containing no 3-hyperedges incident to s .

We give a generalization of this result in Section 1.4.6, for which we will provide a short proof in the appendix, much shorter than the original proof of B. Fleiner of Theorem 10.

Augmentation of global edge-connectivity by attaching stars

By **attaching a star** of degree d to a graph G , we mean adding a new vertex and connecting it to some vertices of G so that the degree of the new vertex becomes d . The problem of this section is the **global edge-connectivity augmentation problem of a graph by attaching stars** that can be defined as follows: *Given a graph $G = (V, E)$ and integers k, d_1, \dots, d_p , decide whether it is possible to attach p stars to G with degrees d_1, \dots, d_p to have a k -edge-connected graph in V .*

Note that, by taking each d_i to be equal to 2, we get as a special case the global edge-connectivity augmentation problem in a graph.

The solution of the problem of this section is given in the following theorem of B. Fleiner which is implied by Theorems 1 and 10. It might be advantageous to notice that condition (1.20) and the fact that $k - d(X)$ is symmetric skew-supermodular guarantees that the minimal extension can be made by $\sum_{j=1}^p d_j$ edges, while condition (1.21) allows us to find the suitable detachment.

Theorem 11 (B. Fleiner [14]). *A graph $G = (V, E)$ can be made k -edge-connected ($k \geq 2$) by attaching p stars with degrees d_1, \dots, d_p ($d_i \geq 2 \forall i$) if and only if*

$$\sum_{X \in \mathcal{X}} (k - d(X)) \leq \sum_{j=1}^p d_j \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.20)$$

$$k - \sum_{j=1}^p \lfloor \frac{d_j}{2} \rfloor \leq \lambda(G). \quad (1.21)$$

Note that, for $d_i = 2 \forall i$, (1.21) is satisfied by (1.20). Indeed, for $X \subset V$ with $d(X) = \lambda(X)$, by (1.20) for $\{X, V - X\}$, $(k - d(X)) + (k - d(V - X)) \leq 2p$, so by the symmetry of $d(X)$, $k - p \leq d(X) = \lambda(G)$, that is, (1.21) is satisfied. Hence Theorem 11 implies Theorem 3.

1.4.6 Local edge-connectivity II

In this section we generalize the results of the previous section. First we extend the Theorem of B. Fleiner to the case when local edge-connectivity is involved, and then we solve the problem where we wish to attach stars of given degree to a graph so that the local edge-connectivity be greater than or equal to a given requirement for each pair of vertices.

Detachments preserving local edge-connectivity

Recall that $G = (V + s, E)$ is \mathbf{r} -edge-connected if $\lambda_G(u, v) \geq r(u, v) \forall u, v \in V$, where $r : V \times V \rightarrow \mathbb{Z}_+$ is a symmetric requirement function. In the next result we characterize graphs that have an \mathbf{r} -edge-connected $f(s)$ -detachment.

Note that, by taking \mathbf{r} to be equal to k for each pair of vertices, we get a characterization of graphs having a k -edge-connected $f(s)$ -detachment.

Theorem 12 (Jordán, Szigeti [24]). *Let $G = (V + s, E)$ be a graph, \mathbf{r} a symmetric requirement function on V with $r(u, v) \geq 2 \forall u, v \in V$ and $f(s) = (d_1, \dots, d_p)$ a degree specification for s with $d_i \geq 2 \forall i$. Let $\varphi = \sum_1^p \lfloor \frac{d_i}{2} \rfloor$. Then there exists an \mathbf{r} -edge-connected $f(s)$ -detachment of G if and only if*

$$G \text{ is } \mathbf{r}\text{-edge-connected}, \quad (1.22)$$

$$G - s \text{ is } (\mathbf{r} - \varphi)\text{-edge-connected}. \quad (1.23)$$

The special case of Theorem 12, when $r(u, v) = k \forall u, v \in V$, is Theorem 10, and when $r(u, v) = \lambda_G(u, v) \forall u, v \in V$, provides a characterization of the existence of an $f(s)$ -detachment that preserves local edge-connectivities, while when at most one d_i is odd, is equivalent to Theorem 4(a).

We will provide a short proof of the above theorem in Section 1.8.1. It will use Theorem 4(a) so it does not provide a new proof for Theorem 4(a). Note that it provides a short proof for Theorem 10.

Augmentation of local edge-connectivity by attaching stars

In this section we solve the following **global edge-connectivity augmentation problem of a graph by attaching stars**: *Given a graph $G = (V, E)$, a symmetric requirement function $r : V \times V \rightarrow \mathbb{Z}_+$ and integers d_1, \dots, d_p , decide whether it is possible to attach p stars to G with degrees d_1, \dots, d_p , so as to obtain an \mathbf{r} -edge-connected graph in V .*

Note that, by taking \mathbf{r} to be equal to k for each pair of vertices, we get as a special case the global edge-connectivity augmentation problem of a graph by attaching stars, and by taking each d_i to be equal to 2 but \mathbf{r} being arbitrary, we get the local edge-connectivity augmentation problem.

Recall that $R(X) = \max\{r(x, y) : x \in X, y \in V - X\}$ and that the function $R(X) - d(X)$ is symmetric skew-supermodular. Then Theorems 1 and 12 imply at once the following theorem. As for the global case, we may notice that condition (1.24) guarantees that the minimal extension can be made by $\sum_{j=1}^p d_j$ edges, while condition (1.25) allows us to find the suitable detachment.

Theorem 13 (Jordán, Szigeti [24]). *Let $G = (V, E)$ be a graph, $r : V \times V \rightarrow \mathbb{Z}_+$ a symmetric requirement function with $r(u, v) \geq 2 \forall u, v \in V$. Then G can be made \mathbf{r} -edge-connected by attaching p stars with degrees d_1, \dots, d_p ($d_i \geq 2 \forall i$) if and only if*

$$\sum_{X \in \mathcal{X}} (R(X) - d(X)) \leq \sum_{j=1}^p d_j \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.24)$$

$$r(u, v) - \sum_{j=1}^p \lfloor \frac{d_j}{2} \rfloor \leq \lambda_G(u, v) \quad \forall u, v \in V. \quad (1.25)$$

Note that, if $r(u, v) = k \forall u, v \in V$, then (1.24) and (1.25) are equivalent to (1.20) and (1.21), so Theorem 13 implies Theorem 11, and if $d_i = 2 \forall i$, then (1.24) implies (1.25) (that can be shown similarly, as it was shown for the global case) and hence Theorem 13 implies Theorem 5. Hence, our result is a common generalization of B. Fleiner's theorem on global edge-connectivity augmentation by attaching stars and Frank's theorem on local edge-connectivity augmentation by adding edges.

1.4.7 Global edge-connectivity with partition constraints

The aim of this section is to present the solution of the problem of global edge-connectivity augmentation in bipartite graphs. In fact, we consider a more general setting, namely the **global edge-connectivity augmentation problem of a graph with partition constraints**: *we want to make an arbitrary graph k -edge-connected by adding a minimum number of new edges between different members of a given partition of the vertex set.*

Splitting off preserving global edge-connectivity with partition constraints

Let $G = (V + s, E)$ be a graph with a specified vertex s of even degree, and $\mathcal{P} = \{P_1, \dots, P_r\}$ a partition of V (resp. $\delta(s)$). If we have a partition of V , then obviously we may define a partition of $\delta(s)$, hence the second form is more general than the first one. A splitting off $\{su, sv\}$ is called **\mathcal{P} -allowed** if u and v (resp. su and sv) belong to different members of \mathcal{P} . A k -admissible \mathcal{P} -allowed splitting off will be called **allowed**. We wish to characterize graphs and partitions for which there exists a complete allowed splitting off, that

is, we are interested in a complete splitting off, that is, at the same time \mathcal{P} -allowed and k -admissible. Notice that a complete \mathcal{P} -allowed splitting off exists if and only if $d(s, P_i) \leq \frac{d(s)}{2} \forall 1 \leq i \leq r$, while, as it is already known for us, a complete k -admissible splitting off exists if and only if G is k -edge-connected in V . Are these conditions together sufficient to have an allowed complete splitting off? We will answer this question in this section.

To show the difficulties of our problem, suppose we wish to make 3-edge-connected a 4-cycle C_4 . Clearly, this can be done by adding two edges. Note that the optimal solution is unique, we have to transform the graph into a K_4 . Now suppose that we have an additional condition, we must maintain the bipartiteness of C_4 . Then this solution is not feasible any more. In this case we have to add 3 edges. Let us reformulate this difficulty in terms of splitting off. Let G be the graph obtained from C_4 by adding a new vertex s and connecting s to all the four vertices. Let \mathcal{P} be the bipartition of C_4 . Then G admits no complete 3-admissible \mathcal{P} -allowed splitting off. The essential properties of this example are kept in a more general structure called C_4 -obstacle, defined as follows.

A partition $\{A_1, A_2, A_3, A_4\}$ of V is called a **C_4 -obstacle** of G if k is odd and

$$d(A_i) = k \quad \forall 1 \leq i \leq 4, \quad (1.26)$$

$$d(A_i, A_{i+2}) = 0 \quad \forall 1 \leq i \leq 2, \quad (1.27)$$

$$|P_l| = \frac{d(s)}{2} \quad \exists 1 \leq l \leq r, \quad (1.28)$$

$$\delta(A_j \cup A_{j+2}) \cap \delta(s) = P_l \quad \exists 1 \leq j \leq 2. \quad (1.29)$$

Another difficulty may turn up if the partition contains more than two members. Suppose we wish to make 3-edge-connected a 6-cycle C_6 . Clearly, this can be done by adding three edges. Note that, though the optimal solution is not unique, we have to add at least one diagonal edge. Now suppose that we have the additional condition that we must maintain the non-adjacency of the opposite vertices of C_6 . Then this solution is not feasible any more. In this case we have to add 4 edges. Let us reformulate this difficulty in terms of splitting off. Let G be the graph obtained from C_6 by adding a new vertex s and connecting s to all the six vertices. Let \mathcal{P} the 3-partition of C_6 where the sets consist of the opposite vertices of C_6 . Then G admits no complete 3-admissible \mathcal{P} -allowed splitting off. As before, the essential properties of this example are kept in a more general structure called C_6 -obstacle, defined as follows.

A partition $\{A_1, \dots, A_6\}$ of V is called a **C_6 -obstacle** of G if k is odd and

$$d(A_i) = k \quad \forall 1 \leq i \leq 6, \quad (1.30)$$

$$d(A_i, A_{i+1}) = \frac{k-1}{2} \quad \forall 1 \leq i \leq 6, (A_7 = A_1) \quad (1.31)$$

$$d(s, A_i) = 1 \quad \forall 1 \leq i \leq 6, \quad (1.32)$$

$$\delta(A_j \cup A_{j+3}) \cap \delta(s) = P_{l_j} \quad \forall 1 \leq j \leq 3, \exists 1 \leq l_j \leq r. \quad (1.33)$$

We must emphasize that these difficulties may exist only if the target edge-connectivity is odd, and if one of them exists, then no complete allowed splitting off may exist. Now we are in a position to provide a characterization of the existence of a complete allowed splitting off.

Theorem 14 (Bang-Jensen, Gabow, Jordán, Szigeti [2]). *Let $G = (V + s, E)$ be a graph with $d(s)$ even, $2 \leq k \in \mathbb{Z}$, and $\mathcal{P} = \{P_1, \dots, P_r\}$ a partition of V . Then there exists a complete k -admissible \mathcal{P} -allowed splitting off at s if and only if*

$$G \text{ is } k\text{-edge-connected in } V, \quad (1.34)$$

$$d(s, P_i) \leq \frac{d(s)}{2} \quad \forall 1 \leq i \leq r, \quad (1.35)$$

$$G \text{ contains no } C_4\text{- or } C_6\text{-obstacle.} \quad (1.36)$$

In the special case, when each element of \mathcal{P} is a singleton, no C_4 or C_6 -obstacle can exist, thus Theorem 14 implies Theorem 2(a), while when $|\mathcal{P}| = 2$, it provides (with Theorem 1) a solution for the problem of global edge-connectivity augmentation in bipartite graphs. We may observe that in this case no C_6 -obstacle can exist.

The following result from [35] is a slight generalization of Theorem 14. The motivation of this form is that it allows us to contract tight sets and hence it enables us to simplify the proof that will be presented in Section 1.8.2.

Theorem 15 (Szigeti [35]). *Let $G = (V + s, E)$ be a graph with $d(s)$ even, $2 \leq k \in \mathbb{Z}$, and $\mathcal{P} = \{P_1, \dots, P_r\}$ a partition of $\delta(s)$. Then there exists a complete k -admissible \mathcal{P} -allowed splitting off at s if and only if (1.34), (1.36) and the following condition are satisfied.*

$$|P_i| \leq \frac{d(s)}{2} \quad \forall 1 \leq i \leq r. \quad (1.37)$$

Augmentation of global edge-connectivity with partition constraints

We present a more precise reformulation of the problem of this section: *Given a graph $G = (V, E)$, an integer k and a partition $\mathcal{P} = \{P_1, \dots, P_r\}$ of V , what is the minimum number $\mathbf{OPT}_{\mathcal{P}}^k$ of \mathcal{P} -allowed edges whose addition results in a k -edge-connected graph?*

Note that, by taking the partition of all singletons in this problem, we get as a special case the global edge-connectivity augmentation problem in graphs.

The following theorem answers this problem. Let $\Phi := \max\{\alpha, \beta_1, \dots, \beta_r\}$ where

$$\alpha := \max\left\{\left\lceil \sum_{X \in \mathcal{X}} \frac{k - d(X)}{2} \right\rceil : \mathcal{X} \in \mathcal{S}(V)\right\},$$

$$\beta_j := \max\left\{\sum_{Y \in \mathcal{Y}} (k - d(Y)) : \mathcal{Y} \in \mathcal{S}(P_j)\right\} \quad \forall 1 \leq j \leq r.$$

It is crucial to point out that Φ is a lower bound for the optimal value, that is,

$$OPT_P^k \geq \Phi. \quad (1.38)$$

Indeed, by adding an edge to G we may decrease the deficiency $k - d(Z)$ of at most two sets in \mathcal{X} and of at most one set in each \mathcal{Y} , and hence we can decrease Φ by at most one.

We will have equality in (1.38), unless G contains one of the following two configurations. These are the structures that force us to have a C_4 - or C_6 -obstacle in any optimal extension.

A partition $\{A_1, A_2, A_3, A_4\}$ of V is called a **C_4 -configuration** of G if k is odd and

$$d(A_i) < k \quad \forall 1 \leq i \leq 4, \quad (1.39)$$

$$d(A_i, A_{i+2}) = 0 \quad \forall 1 \leq i \leq 2, \quad (1.40)$$

$$\sum_{X \in \mathcal{X}_i} (k - d(X)) = k - d(A_i) \quad \exists \mathcal{X}_i \in \mathcal{S}(A_i) \quad \forall 1 \leq i \leq 4, \quad (1.41)$$

$$\mathcal{X}_j \cup \mathcal{X}_{j+2} \in \mathcal{S}(P_l) \quad \exists 1 \leq l \leq r \quad \exists 1 \leq j \leq 2, \quad (1.42)$$

$$k - d(A_i) + k - d(A_{i+2}) = \Phi \quad \forall 1 \leq i \leq 2. \quad (1.43)$$

A partition $\{A_1, A_2, \dots, A_6\}$ of V is called a **C_6 -configuration** of G if k is odd and

$$d(A_i) = k - 1 \quad \forall 1 \leq i \leq 6, \quad (1.44)$$

$$d(A_i, A_{i+1}) = \frac{k-1}{2} \quad \forall 1 \leq i \leq 6, (A_7 = A_1) \quad (1.45)$$

$$\Phi = 3 \quad \forall 1 \leq i \leq 6, \quad (1.46)$$

$$d(A'_i) = k - 1 \quad \exists 1 \leq j_1, j_2, j_3 \leq r \quad \forall 1 \leq i \leq 6 \quad (1.47)$$

$$\exists A'_i \subseteq A_i \cap P_{j_{i-3 \lfloor \frac{i-1}{3} \rfloor}},$$

where the j_i 's must be different in (1.47).

Since these configurations force us to have an obstacle in any optimal extension, in which case there exists no complete allowed splitting off in the extended graph, the existence of a configuration implies that the optimal solution must contain at least $\Phi + 1$ edges. We show that this can be achieved. Using Theorems 1 and 14 we can prove with some effort the following result.

Theorem 16 (Bang-Jensen, Gabow, Jordán, Szigeti [2]). *Let $G = (V, E)$ be a connected graph, $\mathcal{P} = \{P_1, \dots, P_r\}$ a partition of V and $k \geq 2$. Then G can be made k -edge-connected by adding Φ \mathcal{P} -allowed edges (that is $OPT_{\mathcal{P}}^k = \Phi$) unless G contains a C_4 - or C_6 -configuration when we need one more edge (that is $OPT_{\mathcal{P}}^k = \Phi + 1$).*

As the global edge-connectivity augmentation problem *without partition constraints* can be considered as one with partition constraints where each element of \mathcal{P} is a singleton, in which case no C_4 - or C_6 -configuration can exist and $\Phi = \alpha$, Theorem 16 implies Theorem 3. We note that if we want to augment the global edge-connectivity of a *bipartite graph* then $|\mathcal{P}| = 2$ and hence a C_6 -configuration can not exist.

1.5 Hypergraphs

In this section we wish to present problems on edge-connectivity augmentation in hypergraphs and their solutions. We continue with the same structure as before: each subsection is devoted to one problem, and it is divided in two parts; results on splitting off and then the minimax result on augmentation. We mention that some of the problems that were easy in graphs turn out to be already too difficult in hypergraphs.

1.5.1 Augmentation of global edge-connectivity in a hypergraph by adding graph edges

In this section we consider the problem of **global edge-connectivity augmentation in hypergraphs by adding graph edges**, that can be formulated as follows: *Given a hypergraph \mathcal{G} and $k \in \mathbb{Z}^+$, what is the minimum number of graph edges (hyperedges of size two) whose addition results in a k -edge-connected hypergraph?*

Of course, if the hypergraph is in fact a graph, then the problem reduces to the global edge-connectivity augmentation problem in graphs.

One of the main difficulties with hypergraphs that we have to handle the case when $k = 1$. Why is it so? Because by deleting a hyperedge the number of connected components may increase by a large value. This discussion shows the necessity of condition (1.49).

The corresponding splitting off result is the following.

Theorem 17 (Bang-Jensen, Jackson [3]). *Let $\mathcal{G} = (V + s, \mathcal{E})$ be a hypergraph and $\gamma, k \in \mathbb{Z}^+$ so that $d(s) = 2\gamma$ and each edge incident to s is of size two. Then there is a complete k -admissible splitting off at s if and only if*

$$\mathcal{G} \text{ is } k\text{-edge-connected in } V, \quad (1.48)$$

$$c(\mathcal{G} - s - \mathcal{H}) - 1 \leq \gamma \quad \forall \mathcal{H} \subseteq \mathcal{E}, |\mathcal{H}| \leq k - 1. \quad (1.49)$$

If \mathcal{G} is a graph, then the second condition of the above theorem is satisfied (after deleting $k - 1$ edges in a k -edge-connected graph, the graph remains connected) and hence Theorem 17 implies Theorem 2(a).

Theorems 1 and 17 imply the following theorem that solves the problem of this section. We notice that (1.50) and the fact that the function $k - d_{\mathcal{H}}(X)$ is symmetric skew-supermodular guarantees that the optimal extension (with respect to k) can be made by 2γ edges and (1.51) allows us to find the complete k -admissible splitting off.

Theorem 18 (Bang-Jensen, Jackson [3]). *Let \mathcal{G} be a hypergraph and $k \in \mathbb{Z}^+$. Then \mathcal{G} can be made k -edge-connected by adding at most γ new edges (hyperedges of size two) if and only if*

$$\sum_{X \in \mathcal{X}} (k - d_{\mathcal{G}}(X)) \leq 2\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.50)$$

$$c(\mathcal{G} - \mathcal{H}) - 1 \leq \gamma \quad \forall \mathcal{H} \subseteq E(\mathcal{G}), |\mathcal{H}| \leq k - 1. \quad (1.51)$$

As above, if \mathcal{G} is a graph, then (1.51) is implied by (1.50) and hence Theorem 18 implies Theorem 3.

1.5.2 Augmentation of global edge-connectivity of a hypergraph by adding a uniform hypergraph

As a natural generalization of the problem of the previous section, one can consider the **global edge-connectivity augmentation problem of a hypergraph by adding hyperedges of the same size**, namely: *Given a hypergraph \mathcal{G} and $k, r \in \mathbb{Z}^+$, what is the minimum number of hyperedges of size r whose addition results in a k -edge-connected hypergraph?*

Since the function $k - d_{\mathcal{H}}(X)$ is symmetric crossing supermodular, the problem of covering a symmetric crossing supermodular function by a uniform hypergraph contains this problem as a special case, thus Theorem 26 implies the following theorem. We mention that it is not easy to see that (1.54) implies (1.65) when $p(X) = k - d_{\mathcal{G}}(X)$.

Theorem 19 (T. Király [26]). *Let $\mathcal{G} = (V, \mathcal{E})$ be a hypergraph and $k, r \in \mathbb{Z}_+$, $r \leq |V|$. Then \mathcal{G} can be made k -edge-connected by adding at most γ new hyperedges of size r if and only if*

$$\sum_{X \in \mathcal{X}} (k - d_{\mathcal{G}}(X)) \leq r\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.52)$$

$$k - d_{\mathcal{G}}(X) \leq \gamma \quad \forall X \subseteq V, \quad (1.53)$$

$$c(\mathcal{G} - \mathcal{H}) - 1 \leq (r - 1)\gamma \quad \forall \mathcal{H} \subseteq \mathcal{E}, |\mathcal{H}| = k - 1. \quad (1.54)$$

The special case when we want to augment the edge-connectivity from k to $k + 1$ was solved earlier in Fleiner, Jordán [13]. Note that, when $r = 2$, then (1.52) and (1.54) reduce to (1.50) and (1.51), and (1.53) is implied by (1.52), so Theorem 19 implies Theorem 18.

1.5.3 Augmentation of local edge-connectivity of a hypergraph by adding graph edges

This section is devoted to the following problem.

HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH

Instance: A hypergraph \mathcal{H} on V , a symmetric requirement function $r(u, v) \in \mathbb{Z} \forall u, v \in V$, and $\gamma \in \mathbb{Z}_+$.

Question: Does there exist a graph $G = (V, E)$ with at most γ edges so that $\lambda_{\mathcal{H}+G}(u, v) \geq r(u, v) \forall u, v \in V$?

The following theorem shows that this problem is already too complicated.

Theorem 20 (Cosh, Jackson, Z. Király [11]). *The problem HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH is NP-complete.*

We mention that the above problem remains NP-complete if the hypergraph contains only just one hyperedge of size greater than 2.

We remark that the special case of the problem HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH for 2-3 hypergraphs is tractable. This is because of the fact that a 2-3 hypergraph can be transformed into a graph with the same local edge-connectivities and vice versa. Thus Theorem 12 implies the following.

Theorem 21 (Jordán, Szigeti [24]). *Let r be a requirement function with $2 \leq r(u, v) \in \mathbb{Z} \forall u, v \in V$. A 2-3 hypergraph $G = (V, E)$ can be made \mathbf{r} -edge-connected by adding γ edges and γ' 3-hyperedges if and only if*

$$\sum_{X \in \mathcal{X}} (R(X) - d(X)) \leq 2\gamma + 3\gamma' \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.55)$$

$$G \text{ is } \mathbf{r} - (\gamma + \gamma')\text{-edge-connected.} \quad (1.56)$$

1.5.4 Bipartite constrained augmentation of local edge-connectivity of a hypergraph by adding graph edges

In this section we consider a restricted version of the main problem of the preceding section.

BIPARTITION CONSTRAINED HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH

Instance: A hypergraph \mathcal{H} on V , a bipartition $\{A, B\}$ of V , a symmetric requirement function $r(u, v) \in \mathbb{Z} \forall u, v \in V$, and $\gamma \in \mathbb{Z}_+$.

Question: Does there exist a bipartite graph $G = (A, B; E)$ with colour classes A and B with at most γ edges so that $\lambda_{\mathcal{H}+G}(u, v) \geq r(u, v) \forall u, v \in V$?

By checking the proof of Theorem 20 given in [11] we may observe that it also provides the NP-completeness of the above problem.

Theorem 22. *The problem BIPARTITION CONSTRAINED HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH is NP-complete.*

1.5.5 Augmentation of local edge-connectivity of a hypergraph by adding hyperedges

In this section we want to augment a hypergraph by adding hyperedges to satisfy local edge-connectivity requirements. The problem of minimizing the number of hyperedges is trivial since we can add the whole vertex set as many times as needed. What we want to minimize is the total size of the hypergraph to be added, that is, the sum of the sizes of the hyperedges. More precisely, the **local edge connectivity augmentation problem of a hypergraph by adding hyperedges** is the following: *Given a hypergraph \mathcal{G} and a symmetric requirement function $r : V \times V \rightarrow \mathbb{Z}_+$, what is the minimum total size $\sum_{H \in \mathcal{H}} |H|$ of new hyperedges $H \in \mathcal{H}$ whose addition results in an r -edge-connected hypergraph?*

Recall that $R(X) = \max\{r(x, y) : x \in X, y \in V - X\}$ and that the function $R(X) - d_{\mathcal{G}}(X)$ is symmetric skew-supermodular. Then the problem of covering a symmetric skew-supermodular function by a hypergraph contains this problem as a special case thus Theorem 31 implies the following theorem.

Theorem 23 (Szigeti [31]). *Let \mathcal{G} be a hypergraph on V and $r(u, v) \in \mathbb{Z}_+ \forall u, v \in V$ a symmetric requirement function. Then there exists a hypergraph \mathcal{H} on V with $\sum_{H \in \mathcal{H}} |H| \leq \gamma$ so that $\lambda_{\mathcal{G}+\mathcal{H}}(u, v) \geq r(u, v) \forall u, v \in V$ if and only if*

$$\sum_{X \in \mathcal{X}} (R(X) - d_{\mathcal{G}}(X)) \leq \gamma \quad \forall \mathcal{X} \in \mathcal{S}(V). \tag{1.57}$$

1.6 Abstract forms

In this section we present results on "connectivity" set functions, that generalize the results presented in the previous sections. We start with a generalization of splitting off. Let $p : 2^V \rightarrow \mathbb{Z}$ be a function and $H = (V + s, E)$ a graph so that each edge is incident to s . By a **complete hypergraph splitting off** we mean a hypergraph \mathcal{H} on V such that \mathcal{H} covers p and the following is satisfied:

$$d_{\mathcal{H}}(v) = d_H(v) \quad \forall v \in V. \tag{1.58}$$

In other words, it is a degree constrained hypergraph that covers p . Note that there is no restriction on the size of the hyperedges. If we pose the restriction that each hyperedge must be of size at most two, then we are back to the definition of the usual complete splitting off.

1.6.1 Symmetric crossing supermodular functions

The problem of covering a symmetric crossing supermodular function can be considered as a generalization of the global edge-connectivity augmentation

problem. In this section we present the result of Benczúr and Frank on covering a symmetric crossing supermodular function by a graph (that generalizes Theorem 17) and then its generalization due to T. Király on covering such a function by an r -uniform hypergraph (that also generalizes Theorem 19).

Complete uniform hypergraph splitting off

The splitting off results that will solve (together with Theorem 1) the above mentioned problems are the following. We call the attention to the fact that in Theorem 24 a hyperedge may contain the same vertex many times and the size of the hyperedge is meant by multiplicities.

A partition $\{V_1, \dots, V_l\}$ of V is called an **l -partition**. Let $r \in \mathbb{Z}_+$, $H = (V + s, E)$ a graph so that each edge is incident to s and r divides $d_H(s)$ and $p : 2^V \rightarrow \mathbb{Z}_+$ a set function. An l -partition is called **p -full** if $l > r$ and $p(\bigcup_{i \in I} V_i) > 0 \forall \emptyset \neq I \subset \{1, \dots, l\}$. A p -full partition is called **deficient** if $\frac{l-1}{r-1} > \frac{d_H(s)}{r}$. We say that the set function p is **positively crossing supermodular** if (1.10) is satisfied if $X, Y \subseteq V$ are crossing with $p(X), p(Y) > 0$.

Theorem 24 (T. Király [26]). *Let $p : 2^V \rightarrow \mathbb{Z}_+$ be a symmetric, positively crossing supermodular set function, $r \geq 2$ an integer, and $H = (V + s, E)$ a graph so that each edge is incident to s with r divides $d_H(s)$. Then there exists a complete r -uniform hypergraph splitting off if and only if*

$$\min\{d_H(X), d_H(s)/r\} \geq p(X) \quad \forall X \subseteq V, \quad (1.59)$$

$$\text{There are no deficient partitions.} \quad (1.60)$$

The special case when $r = 2$ was proved earlier by Benczúr and Frank [6].

Covering a symmetric crossing supermodular function by a graph

As a generalization of the global edge-connectivity augmentation in hypergraphs by adding graph edges, Benczúr and Frank considered the problem of **covering a symmetric crossing supermodular function by a graph**, namely: *Given a symmetric, positively crossing supermodular set function p , what is the minimum number of edges that cover p ?*

As a generalization of Theorem 18 they proved in [6] the following result by applying Theorem 1 and their splitting off result which is the $r = 2$ special case of Theorem 24.

Theorem 25 (Benczúr, Frank [6]). *Let $p : 2^V \rightarrow \mathbb{Z}_+$ be a symmetric, positively crossing supermodular set function. Then there exists a graph on V with γ edges that covers p if and only if*

$$\sum_{X \in \mathcal{X}} p(X) \leq 2\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V), \quad (1.61)$$

$$l - 1 \leq \gamma \quad \text{if a } p\text{-full } l\text{-partition exists.} \quad (1.62)$$

For a hypergraph \mathcal{G} , the function $k - d_{\mathcal{G}}(X)$ is symmetric, positively crossing supermodular, thus Theorem 25 implies Theorem 18.

Covering a symmetric crossing supermodular function by a uniform hypergraph

As a generalization of the problems of covering a symmetric crossing supermodular function by graph edges and of the global edge-connectivity augmentation in hypergraphs by r -uniform hyperedges, T. Király considered the problem of **covering a symmetric crossing supermodular function by an r -uniform hypergraph**, namely: *Given a symmetric, positively crossing supermodular set function p and $r \in \mathbb{Z}^+$, what is the minimum number of hyperedges of size r that cover p ?*

As a generalization of Theorems 25 and 19 he obtained in [26] the following result by applying Theorems 1 and 24.

Theorem 26 (T. Király [26]). *Let $p : 2^V \rightarrow \mathbb{Z}_+$ be a symmetric, positively crossing supermodular set function, $2 \leq r \leq |V|$ an integer. Then there exists an r -uniform hypergraph on V with γ hyperedges that covers p if and only if*

$$\sum_{X \in \mathcal{X}} p(X) \leq r\gamma \quad \forall \mathcal{X} \in \mathcal{S}(V), \tag{1.63}$$

$$p(X) \leq \gamma \quad \forall X \subseteq V, \tag{1.64}$$

$$l - 1 \leq (r - 1)\gamma \quad \text{if a } p\text{-full } l\text{-partition exists.} \tag{1.65}$$

Note that Theorem 26 implies Theorem 25 (when $r = 2$) and Theorem 19 (when the function is the deficiency function of a hypergraph).

1.6.2 Symmetric skew-supermodular functions

The problem of covering a symmetric skew-supermodular function can be considered as a common generalization of many of the preceding problems. In this section we provide a nice proof of the NP-completeness of the problem of covering a symmetric skew-supermodular function by a graph (which we know already since it generalizes the problem of local edge-connectivity augmentation of a hypergraph by graph edges) while the problem of covering such a function by a hypergraph is solvable in a certain sense.

Covering a symmetric skew-supermodular function by a graph

The problem of this section can be formulated as follows.

MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH

Instance: A symmetric skew-supermodular function p on V and $\gamma \in \mathbb{Z}^+$.

Question: Does there exist a graph on V with at most γ edges that covers p ?

Note that, as we already mentioned, the NP-complete problem HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH is a special case of the problem MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH and hence this last one is also NP-complete, which we prove here in an elegant way. The proof is due to Z. Király [27] and independently to Nutov [30].

Theorem 27. [27] *The problem MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH is NP-complete.*

Proof. [Z. Király] We reduce 3DM to MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH. Let \mathcal{H} be a 3-uniform hypergraph on V . Let $n := |V|$. Let $p(X) := 1$ if $|X| \in \{1, 2, n-1, n-2\}$, or $X \in \mathcal{H}$ or $V-X \in \mathcal{H}$ and 0 otherwise. It is easy to verify that $p(X)$ is a symmetric skew-supermodular set function. The following completes the proof. *There exists a graph on V with at most $2n/3$ edges that covers p if and only if \mathcal{H} contains a 3-dimensional matching.* Indeed, first suppose that $H_1, \dots, H_{n/3}$ is a 3-dimensional matching. For each H_i , let us choose two edges on $V(H_i)$, and let F be the union of these edges. Then $|F| = 2n/3$ and, clearly, F covers p . Now suppose that the graph F covers p and $|E(F)| \leq 2n/3$. Let F_1, \dots, F_l be the connected components of F . Since F covers p , $|V(F_i)| \geq 3$ for $1 \leq i \leq l$ thus $l \leq n/3$. Then $2n/3 \geq |E(F)| \geq n-l \geq 2n/3$ so $l = n/3$ and $|V(F_i)| = 3$ for $1 \leq i \leq l$. Since F covers p , each F_i belongs to \mathcal{H} , that is, F_1, \dots, F_l is a 3-dimensional matching. \square

By the following theorem, which is an easy corollary of Theorem 31, MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH can be solved if $p(X)$ is even for every $X \subseteq V$.

Theorem 28 (Szigeti [31]). *Let p be a symmetric, skew-supermodular, "even integer" valued function on V . Then there exists a graph on V with at most γ edges that covers p if and only if (1.61) is satisfied.*

Proof. To prove the difficult part of Theorem 28, suppose that (1.61) is satisfied and let $p'(X) = p(X)/2$. Then, by the assumptions on p , p' is a symmetric, skew-supermodular, integer valued function on V . Note that, since p satisfies (1.61), p' satisfies (1.13). Then, by Theorem 31, there exists a hypergraph \mathcal{H} on V so that $\sum_{H \in \mathcal{H}} |H| \leq \gamma$ and \mathcal{H} covers p' , that is, $d_{\mathcal{H}}(X) \geq p'(X)$. Let $G := (V, \cup_{H \in \mathcal{H}} E_H)$ where E_H is an arbitrary cycle on $V(H)$ for each $H \in \mathcal{H}$. Then $|E(G)| = \sum_{H \in \mathcal{H}} |E_H| = \sum_{H \in \mathcal{H}} |H| \leq \gamma$ and $d_G(X) = 2d_{\mathcal{H}}(X) \geq 2p'(X) = p(X)$, that completes the proof. \square

Note that the problem MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH contains as a special case : the

global and local edge-connectivity augmentation in graphs and in hypergraphs, the global edge-connectivity augmentation over symmetric parity families, the node to area edge-connectivity augmentation, and the problem of covering a symmetric crossing supermodular function by a graph.

Covering a symmetric semi-monotone function by a graph

In this section, we mention another special case of the NP-complete problem MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH, that can be solved in polynomial time.

We call a function $P : 2^V \rightarrow \mathbb{Z}$ **semi-monotone** if $P(\emptyset) = P(V) = 0$ and for each set $\emptyset \neq X \neq V$, $0 \leq P(X) \leq P(X')$ either for all $\emptyset \neq X' \subseteq X$ or for all $\emptyset \neq X' \subseteq V - X$. We note that a symmetric semi-monotone function is skew-supermodular.

We consider the problem of **Covering a symmetric semi-monotone function by a graph**: *Given a graph $G = (V, E)$ and a symmetric semi-monotone function P on V , add a minimum number $\mathbf{Opt}(P, G)$ of new edges to G to get a covering of P .*

We have already seen in Section 1.4.4 that the node to area global edge-connectivity augmentation problem is a special case of this problem. On the other hand, any instance of the symmetric semi-monotone function covering problem can easily be formulated as a node to area global edge-connectivity augmentation problem. This discussion establishes the surprising equivalence between two seemingly not closely related problems, showing that these are two alternative models of the same problem. (see [20], [18])

We have to emphasize that this problem in general is NP-complete because the function defined in the proof of Theorem 27 is semi-monotone. However, if the function does not take the value 1, then it can be solved.

Let us define $Q(G) := \max\{\sum_{X \in \mathcal{X}} q(X) : \mathcal{X} \in \mathcal{S}(V)\}$, where $q(X) = P(X) - d_G(X)$. Now we have a lower bound for the optimal value, namely $\mathbf{Opt}(P, G) \geq \lceil \frac{Q(G)}{2} \rceil$. Usually equality will hold, unless the graph contains a special configuration.

Since the above defined function $q(X)$ is symmetric skew-supermodular, Theorem 1 and a suitable splitting result [18] provide the solution of the covering problem.

Theorem 29 (Ishii [20], Grappe and Szigeti [18]). *Let $G = (V, E)$ be a graph and P a symmetric semi-monotone function on V so that $P(X) \neq 1 \forall X \subseteq V$. If G contains no configuration, then $\mathbf{Opt}(P, G) = \lceil \frac{Q(G)}{2} \rceil$, otherwise $\mathbf{Opt}(P, G) = \lceil \frac{Q(G)}{2} \rceil + 1$.*

The definition of the configuration (which is fairly complicated) and a short proof of Theorem 29 can be found in [18]. Notice that Theorem 29 implies Theorem 9.

Complete hypergraph splitting off

For symmetric skew-supermodular functions we have the following splitting off result.

Theorem 30 (Szigeti [31]). *Let $p : 2^V \rightarrow \mathbb{Z}$ be a symmetric skew-supermodular function. Let $H = (V + s, E)$ be a graph so that each edge is incident to s . Then there exists a complete hypergraph splitting off if and only if H covers p .*

T. Király [25] has recently found a very short proof for a slight extension of Theorem 30.

Covering a symmetric skew-supermodular function by a hypergraph

In this section we provide the solution for the problem of **covering a symmetric skew-supermodular function by a hypergraph**, that is, a generalization of the problem of hypergraph local edge-connectivity augmentation by hyperedges, and that can be formulated as follows: *Given a symmetric skew-supermodular function p , what is the minimum total size $\sum_{H \in \mathcal{H}} |H|$ of a hypergraph \mathcal{H} that covers p ?*

Theorems 1 and 30 provide the following generalization of Theorem 23.

Theorem 31 (Szigeti [31]). *Let $p : 2^V \rightarrow \mathbb{Z}$ be a symmetric skew-supermodular function. Then there exists a hypergraph \mathcal{H} on V with $\sum_{H \in \mathcal{H}} |H| \leq \gamma$ so that \mathcal{H} covers p if and only if (1.13) is satisfied.*

Note that only the total size of the hypergraph is guaranteed and no information is available on the size of the hyperedges. This is not a surprise in the light of the NP-completeness of the problem MINIMUM COVER OF A SYMMETRIC, SKEW-SUPERMODULAR FUNCTION BY A GRAPH. However, Bernáth and T. Király [7] have recently observed that the hypergraph \mathcal{H} in Theorem 31 can be chosen so that each edge except one is of size two. They also proved that the hypergraph can be chosen so that each edge is of size k or $k + 1$ for some $2 \leq k \in \mathbb{Z}$.

1.7 Open problems

In this section we pose some open problems related to (more precisely: generalizing of) the problems of this paper.

1.7.1 Graphs

The following open problem, called **node to area local edge-connectivity augmentation** and mentioned in [19], is a natural generalization of the node to area global edge-connectivity augmentation problem: *Given a graph $G = (V, E)$, a family \mathcal{W} of sets $W \subseteq V$, and a requirement function $r : \mathcal{W} \times V \rightarrow \mathbb{Z}_+$, add a minimum number of new edges to G so that the resulting graph contains $r(W, v)$ edge-disjoint paths from any area $W \in \mathcal{W}$ to any vertex $v \notin W$.*

A result on **partition constrained detachment preserving global edge-connectivity** would imply Theorems 15 and 10. To be more precise we can consider the following problem: *Given a graph $G = (V + s, E)$, a partition \mathcal{P} of $\delta(s)$, a degree specification $f(s)$ and a positive integer k , decide whether G has an $f(s)$ -detachment that is k -edge-connected in V and \mathcal{P} -allowed (meaning that for each new vertex s_i , the edges incident to s_i belong to different members of \mathcal{P}).*

A characterization of the existence of a **partition constrained complete splitting off satisfying a requirement function** would imply Theorem 14 and also Theorem 12 as it was observed by Frank [17]. Indeed, for a graph $G = (V + s, E)$, a requirement function $r : V \times V \rightarrow \mathbb{Z}_+$ ($r(u, v) \geq 2 \forall u, v \in V$) and a degree specification $f(s) = (d_1, \dots, d_p)$ ($d_i \geq 2 \forall i$), let G' be obtained from G by adding p new vertices s_1, \dots, s_p and connecting each vertex s_i to s by d_i new edges, let $r'(u, v) := r(u, v)$ if $u, v \in V$ and 2 if $\{u, v\} \cap \{s_1, \dots, s_p\} \neq \emptyset$ and let $\mathcal{P} := \{V, \{s_1, \dots, s_p\}\}$. Then G'' is a complete r' -edge-connected \mathcal{P} -allowed splitting off of G' if and only if G'' is a r -edge-connected $f(s)$ -detachment of G .

Note that the complexity of the **local edge-connectivity augmentation problem in a graph with bipartition constraint** is not known, however, as we have seen, the problem BIPARTITION CONSTRAINED HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH is NP-complete.

A common generalization of the above two problems is the problem of the existence of a **partition constrained detachment satisfying a requirement function**.

1.7.2 Hypergraphs

The following problem of **global edge-connectivity augmentation in hypergraphs with hyperedges of given sizes** is still open: *Given a hypergraph \mathcal{G} and $k, d_1, \dots, d_p \in \mathbb{Z}^+$ decide whether it is possible to add p hyperedges of sizes d_1, \dots, d_p to \mathcal{G} to get a k -edge-connected hypergraph.* More generally we can consider the problem of covering a symmetric crossing supermodular function with hyperedges of given sizes. A solution to these problems would imply Theorems 19 and 24, respectively.

We may also consider the problem of **partition constrained complete splitting off preserving global edge-connectivity in hypergraphs**: *Given a hypergraph $\mathcal{G} = (V + s, E)$ such that \mathcal{G} is k edge-connected in V and no hyperedge of size at least three is incident to s , and a partition \mathcal{P} of $\delta(s)$, decide whether there exists a \mathcal{P} -allowed k -admissible complete splitting off at s . A result on this problem would imply Theorem 14. The bipartite case was done by Cosh [10].*

The following problem of **detachment preserving global edge-connectivity in hypergraphs** is also open: *Given a hypergraph $\mathcal{G} = (V + s, E)$ such that \mathcal{G} is k -edge-connected in V and no hyperedge of size at least three is incident to s , and a degree specification $f(s)$, decide whether there exists an $f(s)$ -detachment such that the resulting hypergraph is k -edge-connected in V . A result on this problem would imply Theorems 10 and 17. The case of local edge-connectivity (being a generalization of the problem HYPERGRAPH LOCAL EDGE-CONNECTIVITY AUGMENTATION BY A GRAPH) is NP-complete.*

1.8 Appendix

1.8.1 A short proof of Theorem 12

The following short proof is from [34]. Note that, by Menger's Theorem, G is r -edge-connected if and only if $h_G^r(X) \geq 0 \forall X \subseteq V$, where $h_G^r(X) := d_G(X) - R(X)$. The following basic property of the function h follows from the facts that $d_G(X)$ satisfies both (1.2) and (1.3) and that $R(X)$ is skew-supermodular, and will be used frequently in this section. For any two subsets $X, Y \subseteq V$ at least one of (1.66) and (1.67) holds. If $X \cup Y = V$ then (1.67) always holds (with equality).

$$h_G^r(X) + h_G^r(Y) \geq h_G^r(X \cap Y) + h_G^r(X \cup Y) + 2d_G(X, Y), \quad (1.66)$$

$$h_G^r(X) + h_G^r(Y) \geq h_G^r(X - Y) + h_G^r(Y - X) + 2\bar{d}_G(X, Y). \quad (1.67)$$

Proof of the necessity of Theorem 12

Let $G' := (V + \{s_1, \dots, s_p\}, E)$ be an \mathbf{r} -edge-connected $f(s)$ -detachment of G at s . By (1.4), applied for $X = \{s_1, \dots, s_p\}$, (1.22) is satisfied since $G'/X = G$. By (1.5), applied for every vertex s_i $1 \leq i \leq p$, (1.23) is satisfied since $G' - X = G - s$. \square

Proof of the sufficiency of Theorem 12

Wlog. $p \geq 2$ and $\varphi \geq 2$. We will use induction on $z(G) := |V| + d_G(s)$. As we already mentioned, (1.22) and (1.23) can be reformulated as (1.68) and (1.69). Note that (1.70) holds.

$$h_G^r(X) \geq 0 \quad \forall X \subseteq V, \quad (1.68)$$

$$h_{G-s}^{r-\varphi}(X) \geq 0 \quad \forall X \subseteq V. \quad (1.69)$$

$$h_{G-s}^{r-\varphi}(X) = h_G^r(X) - d_G(s, X) + \varphi \quad \forall X \subseteq V. \quad (1.70)$$

Lemma 2. *We may assume that*

$$\text{every set } \emptyset \neq X \subset V \text{ with } h_G^r(X) = 0 \text{ is a singleton.} \quad (1.71)$$

Proof. Suppose there exists a set Q with $h_G^r(Q) = 0$ and $|Q| > 1$. Then let $\hat{G} := (\hat{V}, \hat{E})$ be obtained from G by contracting Q into a vertex q and let $\hat{r}(u, v) := r(u, v)$ if $u, v \in \hat{V} - q$, and $\max\{r(w, x) : w \in Q\}$ if $q \in \{u, v\}$ where $x = \{u, v\} - q$. It can be verified easily that $\hat{R}(\hat{X}) = R(X) \forall \hat{X} \subseteq \hat{V}$, so (1.68) and (1.69) are satisfied for \hat{G} and \hat{r} . Since $|Q| > 1$, $z(\hat{G}) < z(G)$ and hence, by induction, \hat{G} has an \hat{r} -edge-connected $f(s)$ -detachment \hat{G}' . We show that the graph G' obtained from \hat{G}' by “blowing up” Q is r -edge-connected and we are done. Let $X' \subseteq V'$. Using that $h_{G'}^r(Q) = h_G^r(Q) = 0$, the skew-submodularity of $h_{G'}^r$ and the fact that if X' and Q are not intersecting then $h_{G'}^r(X') \geq 0$ (because if $X' \subset Q$ then $h_{G'}^r(X') = h_G^r(X) \geq 0$ by (1.68) and if $Q \subseteq X'$ or $Q \cap X' = \emptyset$ then $h_{G'}^r(X') = h_{\hat{G}'}^{\hat{r}}(\hat{X}') \geq 0$ since \hat{G}' is \hat{r} -edge-connected) we get that $h_{G'}^r(X') \geq 0$ as we wanted. \square

For $T \subset \delta_G(s)$, the **T-split** of G is the $(|T|, d_G(s) - |T|)$ -detachment G' of G at s where $\delta_{G'}(s_1) = T$. For $X \subseteq V$, let $e(\mathbf{T}, \mathbf{X}) = |T \cap \delta_G(X)|$.

Lemma 3. *There exists $T \subset \delta_G(s)$ with $|T| = 3$ if $f(s) = (3, 3, \dots, 3)$ and $|T| = 2$ otherwise such that the T -split G' of G satisfies (1.72) and (1.73) where $\mathbf{r}'(u, v) := r(u, v)$ if $u, v \in V$ and 2 otherwise and $V' = V \cup s_1$.*

$$G' \text{ is } \mathbf{r}'\text{-edge-connected in } V', \quad (1.72)$$

$$G' - s \text{ is } (\mathbf{r}' - (\varphi - 1))\text{-edge-connected in } V', \quad (1.73)$$

Proof. Let \mathcal{C} be defined as the minimal sets X with $h_{G-s}^{r-\varphi}(X) = 0$.

Claim. (1.72) and (1.73) are equivalent to

$$h_G^r(X) \geq 2e(T, X) - |T| \quad \forall X \subset V, \quad (1.74)$$

$$e(T, C) \geq 1 \quad \forall C \in \mathcal{C}. \quad (1.75)$$

Proof. (1.72) is satisfied if and only if $0 \leq h_{G'}^{\mathbf{r}'}(X') \forall X' \subset V'$. Since, for $X' \subset V' - s_1$, $h_{G'}^{\mathbf{r}'}(X') = h_G^r(X) \geq 0$, (1.72) is equivalent to $0 \leq h_{G'}^{\mathbf{r}'}(X') \forall X' \subset V'$ containing s_1 which is, by $h_{G'}^{\mathbf{r}'}(X') = h_G^r(X) - e(T, X) + (|T| - e(T, X))$ with $X = X' - s_1$, equivalent to (1.74). (1.73) is satisfied if and only if $0 \leq h_{G'-s}^{\mathbf{r}'-\varphi'}(X) \forall X \subset V'$ not containing s_1 which is, by $h_{G'-s}^{\mathbf{r}'-\varphi'}(X) = h_{G-s}^{r-\varphi}(X) + e(T, X) - 1$, equivalent to (1.75). \square

Claim. The following are true for \mathcal{C} and for all $C \in \mathcal{C}$:

$$\text{the sets in } \mathcal{C} \text{ are pairwise disjoint and } d_G(s, C) \geq \varphi, \quad (1.76)$$

$$|\mathcal{C}| \in \{0, 2, 3\}, \text{ if } |\mathcal{C}| = 3 \text{ then } f(s) = (3, 3, \dots, 3) \text{ and } h_G^r(C) = 0. \quad (1.77)$$

Proof. By the skew-submodularity of $h_{G-s}^{r-\varphi}(X)$, the minimality of the sets in \mathcal{C} , (1.69), (1.70) and (1.68), (1.76) follows. If $X \in \mathcal{C}$, then $V - X$ contains a set $Y \in \mathcal{C}$ so $|\mathcal{C}| \neq 1$. By (1.76) and $d_i \geq 2$, $|\mathcal{C}|\varphi \leq \sum_{C \in \mathcal{C}} d_G(s, C) \leq d_G(s) = \sum_{i=1}^p d_i \leq 3 \sum_{i=1}^p \lfloor \frac{d_i}{2} \rfloor = 3\varphi$ thus $|\mathcal{C}| \leq 3$ and if $|\mathcal{C}| = 3$ then each $d_i = 3$, that is, $f(s) = (3, 3, \dots, 3)$ and $\forall C \in \mathcal{C}$, $d_G(s, C) = \varphi$, so by (1.70), $h_G^r(C) = 0$. \square

By (1.77), either $|\mathcal{C}| = 3$ or $|\mathcal{C}| \in \{0, 2\}$. If $|\mathcal{C}| = \mathbf{3}$, then, by (1.77), $f(s) = (3, 3, \dots, 3)$. By (1.76), there exists $T \subset \delta_G(s)$ with $|T| = 3$ that satisfies (1.75). T also satisfies (1.74). Indeed, by (1.77), (1.71) and (1.76), $d_G(s, X) \geq \varphi e(T, X)$. So, by (1.70), (1.69) and $\varphi \geq 2$, $h_G^r(X) \geq d_G(s, X) - \varphi \geq \varphi(e(T, X) - 1) \geq 2(e(T, X) - 1) \geq 2e(T, X) - |T|$.

From now on $|\mathcal{C}| \in \{\mathbf{0}, \mathbf{2}\}$.

Claim. There exists $T = \{su, sv\}$ that satisfies (1.74) and (1.75).

Proof. If $|\mathcal{C}| = \mathbf{0}$, then (1.75) is redundant, and, by Theorem 4(a), there is a pair $T = \{su, sv\}$ that is λ -admissible which is equivalent to (1.74). If $\mathcal{C} = \{\mathbf{C}_1, \mathbf{C}_2\}$, then, by (1.76), there is a $T = \{su, sv\}$ satisfying (1.75). T also satisfies (1.74). Otherwise, there exists $u, v \in X \subset V$ with $h_G^r(X) \leq 1$. If $C_1 \cup C_2 \subseteq X$, then, by (1.70), (1.69), (1.76), $h_G^r(X) \geq d_G(s, X) - \varphi \geq d_G(s, C_1 \cup C_2) - \varphi \geq 2\varphi - \varphi \geq 2$, contradiction, so wlog. $Y := C_1 - X \neq \emptyset$. Since $C_1 \in \mathcal{C}$, $h_{G-s}^{r-\varphi}(Y) \geq 1$. Let $d := d_G(s, C_1 \cap X)$. Then, by (1.70), $h_G^r(Y) = h_{G-s}^{r-\varphi}(Y) + d_G(s, Y) - \varphi \geq 1 + (d_G(s, C_1) - d) - \varphi = h_G^r(C_1) + 1 - d$. If (1.67) applies for C_1 and X , then, by (1.68), $h_G^r(X) + h_G^r(C_1) \geq h_G^r(X - C_1) + h_G^r(Y) + 2d_G(X, C_1) \geq h_G^r(Y) + 2d \geq h_G^r(C_1) + 1 + d \geq h_G^r(C_1) + 2$, contradiction. So (1.66) applies for C_1 and X and $Z := C_1 \cup X \neq V$. Since $\mathcal{C} = \{C_1, C_2\}$, $h_{G-s}^{r-\varphi}(Z) = h_{G-s}^{r-\varphi}(V - Z) \geq 1$. Then, by (1.70), $h_G^r(Z) = h_{G-s}^{r-\varphi}(Z) + d_G(s, Z) - \varphi \geq 1 + (d_G(s, C_1) + 1) - \varphi = h_G^r(C_1) + 2$, so, by (1.68), $h_G^r(X) + h_G^r(C_1) \geq h_G^r(X \cap C_1) + h_G^r(Z) \geq h_G^r(C_1) + 2$, a contradiction that finishes the proof of the claim. \square

If $f(s) \neq (3, 3, \dots, 3)$, then, by the above claim, we are done. From now on $f(s) = (\mathbf{3}, \mathbf{3}, \dots, \mathbf{3})$. Then $d_G(s) = 3\varphi$.

Claim. T can be extended to $T' \subset \delta_G(s)$ with $|T'| = 3$ such that T' satisfies (1.74).

Proof. First suppose $\Gamma(s) = \{\mathbf{u}, \mathbf{v}\}$. Since $d_G(s) = 3\varphi$ and $\varphi \geq 2$, wlog. $d_G(s, u) \geq \varphi + 1$, thus there exists another copy e' of su . Then, by (1.70) and (1.69), $T' := T \cup e'$ satisfies (1.74). Hence $\Gamma(s) \neq \{\mathbf{u}, \mathbf{v}\}$. Suppose indirect that there exists a minimal set \mathcal{M} of subsets of V such that for

every $z_i \in \Gamma(s) - \{u, v\} \neq \emptyset$ there exists a set $M_i \in \mathcal{M}$ violating (1.74) for $T' := T \cup sz_i$. Then, since T satisfies (1.74) and by (1.71), $e(T', M_i) = 3$ so $\{u, v, z_i\} \subseteq M_i$ and $h_G^r(M_i) \leq 2$. Clearly, $|\mathcal{M}| \geq 1$. By (1.70), (1.69), $h_G^r(M_i) \leq 2$ and $\varphi \geq 2$, we have $|\mathcal{M}| \geq 2$. For $M_i, M_j \in \mathcal{M}$,

$$h_G^r(M_i - M_j) = 0, \text{ (so, by (1.71), } M_i - M_j = z_i, \text{)} \quad (1.78)$$

$$\bar{d}_G(M_i, M_j) = 2, \quad (1.79)$$

$$d_G(z_i, M_i - z_i) \geq 1. \quad (1.80)$$

Indeed, $2 \geq h_G^r(M_i), 2 \geq h_G^r(M_j), h_G^r(M_i \cap M_j) \geq 2e(T, M_i \cap M_j) - |T| \geq 2 \times 2 - 2 = 2$ (by (1.74) and $\{u, v\} \subset M_i \cap M_j$), $h_G^r(M_i \cup M_j) \geq 3$ (by the minimality of \mathcal{M}), so (1.66) cannot be satisfied for M_i and M_j . Then M_i and M_j satisfy (1.67) implying (1.78) and (1.79). Moreover, since $\max\{R(z_i), R(M_i - z_i)\} \geq R(M_i)$ and $\min\{R(z_i), R(M_i - z_i)\} \geq 2$, we have $R(z_i) + R(M_i - z_i) \geq R(M_i) + 2$, thus $2 \leq h_G^r(z_i) + h_G^r(M_i \cap M_j) = h_G^r(z_i) + h_G^r(M_i - z_i) \leq h_G^r(M_i) - 2 + 2d_G(z_i, M_i - z_i) \leq 2d_G(z_i, M_i - z_i)$ implying (1.80).

Case 1 If $\mathcal{M} = \{M_1, M_2\}$. Then, by (1.70), (1.78), (1.69) and (1.79), $3\varphi = d_G(s) = d_G(s, z_1) + d_G(s, z_2) + d_G(s, M_1 \cap M_2) = h_G^r(z_1) - h_{G-s}^{r-\varphi}(z_1) + \varphi + h_G^r(z_2) - h_{G-s}^{r-\varphi}(z_2) + \varphi + d_G(s, M_1 \cap M_2) \leq 2\varphi + 2 \leq 3\varphi$. Thus $h_{G-s}^{r-\varphi}(z_1) = 0$, so $z_1 \in \mathcal{C}$, that is, (1.75) is violated for T , contradiction.

Case 2 If $M_1, M_2, M_3 \in \mathcal{M}$. Then, by (1.80), (1.78), (1.79), $1 \leq d_G(M_3 - z_3, z_3) = d_G(M_1 \cap M_2, z_3) \leq \bar{d}_G(M_1, M_2) - d_G(M_1 \cap M_2, s) \leq 2 - 2 = 0$, contradiction. The proof of the claim is finished. \square

Since T satisfies (1.75), so does T' and the proof of Lemma 3 is complete.

\square

Let G' be the T -split of G from Lemma 3. Let us denote the new vertex of G' of degree $|T|$ by t . Wlog. $d_1 \geq d_2 \geq \dots \geq d_p$. If $d_p = |T|$ then let $f'(s) := (d_1, \dots, d_{p-1})$ otherwise ($|T| = 2, d_1 \geq 4$) let $f'(s) := (d_1 - 2, d_2, \dots, d_p)$. Then $(G', f'(s))$ satisfies (1.72) and (1.73) and $z(G') < z(G)$, so by induction, G' has an \mathbf{r} -edge-connected $f'(s)$ -detachment G'' . Then, in the former case G'' , in the latter case the graph obtained from G'' by identifying s_1 and t , is an \mathbf{r} -edge-connected $f(s)$ -detachment of G . $\square \square \square$

1.8.2 A short proof of Theorem 15

I hope that the reader will find interesting the following shortened proof from [35]. I ask the reader not to be frightened by the technical aspects of the proof and to make figures to help to understand the proofs. In this section we call a set $X \subset V$ **tight** (resp. **dangerous**) if $d(X) = k$ (resp. $d(X) \leq k + 1$). We will abbreviate k -admissible by admissible. In order to have a more convenient notation, $e \in P_j$ will also be denoted by $c(e) = j$.

Preliminaries

The following easy observations are from [16].

Proposition 1. (a) $\{su, sv\}$ is admissible if and only if there is no dangerous set containing u and v .

(b) For any edge su , there exist at most two dangerous sets M_1 and M_2 so that $u \in M_1 \cap M_2$ and $\{v : \{su, sv\} \text{ is not admissible}\} \subseteq M_1 \cup M_2$.

(c) For a tight set T , $\{su, sv\}$ is admissible in G if and only if it is admissible in G/T . \square

The following proposition contains some technical remarks.

Proposition 2. (a) $d(X) - k \geq 2d(s, X) - d(s) \forall X \subset V$ where equality holds if and only if $d(V - X) = k$.

(b) If $k \geq 3$ and $d(X) \leq k + 2$ then $G[X]$ is connected.

(c) If k is odd, X_1, X_2, X_3 are disjoint tight sets, $d(\cup_{i=1}^3 X_i) = k + 2$ and $d(X_1, X_3) = 0$, then $d(X_1, X_2) = d(X_2, X_3) = \frac{k-1}{2}$.

Proof. (a) By (1.34), $d(X) - k = d(V - X) - k + d(s, X) - (d(s) - d(s, X)) \geq 2d(s, X) - d(s)$.

(b) For a set $\emptyset \neq Y \subset X$, by (1.2) and (1.34), $(k + 2) + 2d(Y, X - Y) \geq d(X) + 2d(Y, X - Y) = d(Y) + d(X - Y) \geq k + k \geq k + 3$, and (b) follows.

(c) By (1.2) and (1.34), $\forall i \in \{1, 3\}$, $2k = d(X_2) + d(X_i) = d(X_2 \cup X_i) + 2d(X_2, X_i) \geq k + 2d(X_2, X_i)$, thus, by parity, $2d(X_2, X_i) \leq k - 1$. Then $3k = \sum_{i=1}^3 d(X_i) = d(\cup_{i=1}^3 X_i) + \sum_{i \neq j} 2d(X_i, X_j) \leq (k + 2) + 2(k - 1) + 0 = 3k$, and (c) follows. \square

We present now some important properties of C_4 - and C_6 -obstacles.

Proposition 3. (a) If \mathcal{A} is a C_4 -obstacle, then $d(s, A_i) \geq 1 \forall A_i \in \mathcal{A}$.

(b) If $\{A_1, A_2, A_3, A_4\}$ is a C_4 -obstacle, then for each set $\emptyset \neq X \subseteq A_i$, $d(V - X) \geq k + 2$ with equality only if $d(X) = k$.

(c) If $\{A_1, a_2, a_3, a_4\}$ is a C_4 -obstacle, then for each set X with $X \cap A_1 \neq \emptyset$ and $a_3 \in X$, $d(X) \geq k + 2$.

(d) If $\{A_1, \dots, A_6\}$ is a C_6 -obstacle, then for every allowed pair $\{sx, sy\}$, $G_{x,y}$ contains a C_4 -obstacle.

(e) If $\{A_1, a_2, \dots, a_6\}$ is a C_6 -obstacle and, for a set $X \neq V$, $X \cap A_1 \neq \emptyset$, $y \in X \cap (a_3 \cup a_5)$ and $d(X) \leq k + 2$, then (e1) $d(X) = k + 2$ and (e2) $X \cup A_1$ is the union of three consecutive sets in \mathcal{A} .

Proof. (a) Suppose wlog. $d(s, A_1) = 0$. Then, by (1.27) and (1.26), $d(A_1, A_2) + d(A_1, A_4) = d(A_1) = k$, so, since k is odd, wlog. $d(A_1, A_2) \geq \frac{k+1}{2}$. Then, by (1.34), (1.2) and (1.26), $k \leq d(A_1 \cup A_2) = d(A_1) + d(A_2) - 2d(A_1, A_2) \leq k + k - (k + 1) = k - 1$, contradiction.

(b) Let $j := i + 2$ if $i \leq 2$ and $j := i - 2$ if $i \geq 3$. Then, by (1.34), Proposition 3(a), (1.28) and (1.29), $k + 2 \leq d(X) + 2d(s, A_j) = d(X) - (\frac{d(s)}{2} - d(s, A_j)) + \frac{d(s)}{2} + d(s, A_j) = d(V - X)$ and (b) follows.

(c) If $a_2, a_4 \notin X$, then, by $X \cap A_1 \neq \emptyset$, (1.34), (1.26) and (1.27), $d(X) = d(X \cap A_1) + d(a_3) = k + k \geq k + 2$. If $a_2, a_4 \in X$, then, by Proposition 3(b), $d(X) \geq k + 2$. Otherwise, wlog. $a_2 \notin X$ and $a_4 \in X$. By Proposition 3(b), $d(X \cup A_1) \geq k + 2$. Then, by (1.2) and (1.34), $d(X) \geq d(X \cap A_1) + d(X \cup A_1) - d(A_1) \geq k + (k + 2) - k = k + 2$.

(d) Wlog. $x \in A_1$. By (1.2), (1.30), (1.31), $d(A_i \cup A_{i+1}) = d(A_i) + d(A_{i+1}) - 2d(A_i, A_{i+1}) = k + k - (k - 1) = k + 1$. Then, since $\{sx, sy\}$ is admissible, $y \notin A_2 \cup A_6$ by Proposition 1(a). $\{sx, sy\}$ is allowed so, by (1.33), $y \notin A_4$. Thus wlog. $y \in A_3$. Then $\{A_1 \cup A_2 \cup A_3, A_4, A_5, A_6\}$ is a C_4 -obstacle in $G_{x,y}$.

(e) Let $X^* := X \cup A_1$. By (1.31), $d_{G-s}(X^*) \geq k - 1$ where equality holds if and only if X^* is the union of $2 < l < 6$ consecutive sets in \mathcal{A} . By Proposition 2(a), $d(s, X) \leq 4$, by (1.32), $d(s, A_1) = 1$ and $d(s, V) = 6$ so $X^* \neq V$. By (1.30), $d(A_1) = k$, by $X \cap A_1 \neq \emptyset$ and (1.34), $d(X \cap A_1) \geq k$, so by (1.2), $(k + 2) + k \geq d(X) + d(A_1) \geq d(X \cap A_1) + d(X^*) \geq k + d(X^*)$, so $k + 2 \geq d(X^*)$ and if equality holds then $d(X) = k + 2$. Then, by Proposition 2(b), $G[X^*]$ is connected. Since $d_{G-s}(y, A_1) = 0$, $X' := X - (y \cup A_1) \neq \emptyset$. Then $k + 2 \geq d(X^*) = d(s, X^*) + d_{G-s}(X^*) \geq d(s, y) + d(s, X') + d(s, A_1) + d_{G-s}(X^*) \geq 1 + 1 + 1 + (k - 1)$, so $d(X^*) = k + 2$ and hence $d(X) = k + 2$ (implying (e1)), $d(s, X') = 1$ and $d_{G-s}(X^*) = k - 1$, thus (e2) is satisfied. \square

The following lemma will allow us to easily find an allowed pair. The main difficulty of the proof of Theorem 15 will be to show that there exists an allowed pair whose splitting off creates no C_4 - and no C_6 -obstacle.

Lemma 4. *If G contains no C_4 -obstacle and (1.37) is satisfied then each edge su belongs to an allowed pair.*

Proof. Let $S := \{sv \in E : \{su, sv\} \text{ is admissible}\}$. Suppose su belongs to no allowed pair. Then every $sv \in S$ and su belong to the same P_j . Then, by (1.37), $\frac{d(s)}{2} \geq |P_j| \geq |S| + 1$, so $|S| \leq \frac{d(s)}{2} - 1$ and if equality holds then $\frac{d(s)}{2} = |P_j|$. It also follows, by Proposition 1(b), that there are at most two dangerous sets M_1 and M_2 so that $u \in M_1 \cap M_2$ and $\{v_i : sv_i \in \delta(s) - S\} \subseteq M_1 \cup M_2$. In fact there are exactly two, because, by Proposition 2, $d(M_1 \cup M_2) - k \geq 2d(s, M_1 \cup M_2) - d(s) = 2(d(s) - |S|) - d(s) \geq d(s) - 2(\frac{d(s)}{2} - 1) = 2$, and if equality holds then $d(V - M_1 \cup M_2) = k$ and $|S| = \frac{d(s)}{2} - 1$. The following claim provides a contradiction.

Claim. $\{A_1 = M_1 \cap M_2, A_2 = M_1 - M_2, A_3 = V - M_1 \cup M_2, A_4 = M_2 - M_1\}$ forms a C_4 -obstacle.

Proof. Note that $A_i \neq \emptyset$ $1 \leq i \leq 4$ and $\cup_{i=1}^4 A_i = V$. By (1.2), (1.34) and $d(M_1 \cup M_2) \geq k + 2$, $2(k + 1) \geq d(M_1) + d(M_2) = d(A_1) + d(M_1 \cup M_2) + 2d(M_1, M_2) \geq k + (k + 2)$, so $d(A_1) = k$, $d(M_1 \cup M_2) = k + 2$ and hence $d(A_3) = k$ and $\frac{d(s)}{2} = |P_j|$ so (1.28) is satisfied, and $d(A_2, A_4) = d(M_1, M_2) = 0$. By (1.3) and (1.34), $2(k + 1) \geq d(M_1) + d(M_2) = d(A_2) + d(A_4) + 2d(A_1, A_3 + s) \geq k + k + 2d(A_1, A_3) + 2d(A_1, s) \geq 2k + 0 + 2$, so $d(A_2) = d(A_4) = k$, $d(A_1, A_3) = 0$

and $d(s, A_1) = 1$. It also follows that $\delta(A_1 \cup A_3) \cap \delta(s) = P_j$, so (1.26), (1.27) and (1.29) are satisfied. This completes the proof of the claim and also of Lemma 4. \square

Proof of the necessity of Theorem 15

Suppose there exists a graph that has a complete allowed splitting off $\{\{e_i, f_i\} : 1 \leq i \leq \frac{d_G(s)}{2}\}$ and violates (1.37) or (1.36). Choose such a graph G with $d_G(s)$ minimum. For every $1 \leq i \leq \frac{d_G(s)}{2}, 1 \leq j \leq r, |P_j \cap \{e_i, f_i\}| \leq 1$ so (1.37) is satisfied, whence G contains a C_4 - or a C_6 -obstacle. By Proposition 3(a) and (1.32), $d_G(s) \neq 0$. Then, either by (1.28) and (1.29) or by Proposition 3(d), G_{e_1, f_1} contains a C_4 -obstacle, G_{e_1, f_1} has of course a complete allowed splitting off, and $d_{G_{e_1, f_1}}(s) < d_G(s)$, contradiction. \square

Proof of the sufficiency of Theorem 15

Induction on $|V|$. By Proposition 1(c), we may assume that

$$\text{every tight set is a singleton.} \quad (1.81)$$

Wlog. $|P_1|$ is maximum. By Lemma 4, there is an allowed pair $\{e = sx, f = sy\}$ with $sx \in P_1$. If, after splitting off this pair, we get stuck by a C_6 -obstacle, then we can get rid of it by the following lemma.

Lemma 5. *Suppose that $G' := G_{e, f}$ contains a C_6 -obstacle $\mathcal{A} = \{A_1, \dots, A_6\}$. Then there exists an edge $f' = sy'$ such that $\{e, f'\}$ is allowed in G and $G'' := G_{e, f'}$ satisfies (1.34), (1.37) and (1.36).*

Proof. The suitable y' will be chosen as follows: Wlog. $x \in A_1$. Then, by (1.30) and (1.81), $A_j = a_j \forall 2 \leq j \leq 6$. By (1.33), $c(sa_3) \neq c(sa_5)$ so either $c(sa_3) \neq 1$ in which case let $y' := a_3$ or $c(sa_5) \neq 1$ and then let $y' := a_5$. We show that y' will do. For each set X with $x, y' \in X \neq V$, by Proposition 3(e), $k + 2 \leq d_{G'}(X) \leq d_G(X)$, so by Proposition 1(a), $\{e, f'\}$ is admissible in G , thus G'' satisfies (1.34). Since $c(sx) = 1 \neq c(sy')$, G'' satisfies (1.37) and $\{e, f'\}$ is allowed in G . It remains to show that G'' satisfies (1.36). Since $xy \in E(G')$, either Case a: $y \in A_1$ or Case b: wlog. $y \in A_2$. In both cases we suppose indirect that G'' contains a C_4 - (Case 1) or a C_6 -obstacle (Case 2) \mathcal{A}' .

Case a: Wlog. $y' = a_5$ and $x \in A'_1$. Suppose $y' \notin A'_1$. By (1.30), $k + 2 = d_{G'}(A_1) + 2 = d_G(A_1)$. By (1.26) or (1.30), $d_G(A'_1) = k$ and then, by (1.81), $|A'_i| = 1 \forall A'_i \in \mathcal{A}'$. Hence $|V| = 4$ or 6 . But $|V| \geq 6$ because G' contains the C_6 -obstacle \mathcal{A} , so $|V| = 6$. Thus $A'_1 = A_1$ and hence $k = d_{G''}(A'_1) = d_{G'}(A_1) = d_G(A_1)$, contradiction. Thus $y' \in A'_1$ and $d_G(A'_1) = k + 2$. Since $d_{G'}(A'_1) \leq d_G(A'_1)$, $A'_1 \cup A_1$ is the union of three consecutive sets in \mathcal{A} by Proposition 3(e).

Case a1: Then $3 = |V - (A_1 \cup A'_1)| \leq |V - A'_1| = 3$ by (1.81) so $A_1 \subset A'_1$ thus, by (1.31), wlog. $A'_j = a_j$ $2 \leq j \leq 4$. By (1.33) for \mathcal{A} , there is a $w \in A_1$ with $c(sw) = c(sa_4)$ but $w \in A'_1$ and $a_4 \in A'_4$, contradiction by (1.29) for \mathcal{A}' .

Case a2: Then $a_6 \in A'_1$. Wlog. $A'_2 = a_4$ and $A'_3 = a_3$. Then, by (1.33) for \mathcal{A} and \mathcal{A}' , $c(sa_3) = c(sa_6) \neq c(sa_3)$, contradiction, and we are done in Case a.

Case b: Then, by (1.30) and (1.81), $A_1 = a_1$ so $|V| = 6$.

Case b1: Since $d_{G''}(s, y') = 0$, wlog. $x, y' \in A'_1$ and $d_G(A'_1) = k + 2$. Since $d_{G'}(A'_1) \leq d_G(A'_1)$, $d_{G'}(A'_1) = k + 2$ and $A'_1 \cup A_1$ is the union of three consecutive sets in \mathcal{A} by Proposition 3(e). Then $d_{G'}(A'_1) = d_G(A'_1)$ so $y' = a_5$. Thus $A'_1 = \{a_5, a_6, a_1\}$. Wlog. $A'_j = a_j$ $2 \leq j \leq 4$ by (1.31) for \mathcal{A} and (1.27) for \mathcal{A}' . By (1.33) for \mathcal{A} , $c(sa_1) = c(sa_4)$, contradiction by (1.29) for \mathcal{A}' .

Case b2: Note that, by (1.32), $d_G(s, a_1) = d_G(s, a_2) = 2$ and $d_G(s, a_h) = 1$ ($3 \leq h \leq 6$). Then $d_{G''}(s, a_2) = 2$ that is, by (1.32), a contradiction, and we are done in Case b, and the proof of Lemma 5 is complete. \square

The following lemma allows us to get rid of a C_4 -obstacle if during the splitting off we get stuck with one.

Lemma 6. *Suppose that $G' := G_{e,f}$ contains a C_4 -obstacle $\mathcal{A} := \{A_1, A_2, A_3, A_4\}$. Then there exists an allowed pair $e' = sx', f' = sy'$ in G such that $G'' := G_{e',f'}$ satisfies (1.34), (1.37) and (1.36).*

Proof. Wlog. $x \in A_1$. Since $xy \in E(G')$, either Case a: wlog. $y \in A_2$ or Case b: $y \in A_1$. By (1.26) and (1.81), $A_j = a_j$ $\forall 2 \leq j \leq 4$, in Case a: $A_1 = a_1$ so $|V| = 4$ and in Case b: $d_G(A_1) = k + 2$.

Case a: We chose e' and f' as follows. If there exists an edge $g = sa_3$ with $c(g) \neq c(e)$ then let $e' := e, f' := g$. Otherwise, since \mathcal{A} is not a C_4 -obstacle in G , there is an edge $h = sa_1$ with $c(h) \neq c(e)$ and then let $e' := sa_3, f' := h$. We show that this pair will do. Note that $\{x', y'\} = \{a_1, a_3\}$. For each set X with $x', y' \in X \neq V$, either $X = \{x', y'\}$ and then, by (1.26) and (1.27), $d_G(X) = d_G(a_1) + d_G(a_3) = k + k \geq k + 2$ or, by $X \neq V$, $\exists i \in \{2, 4\}$ $X = \{a_1, a_3, a_i\}$ and then, by Proposition 3(b), $k + 2 \leq d_{G'}(X) \leq d_G(X)$. In both cases, by Proposition 1(a), $\{e', f'\}$ is admissible in G , thus G'' satisfies (1.34). Since $c(e') = c(e)$, G'' satisfies (1.37) and $\{e', f'\}$ is allowed in G . It remains to show that G'' satisfies (1.36). Suppose not. Then, since $|V| = 4$, G'' contains a C_4 -obstacle $\mathcal{A}' := \{A'_1, A'_2, A'_3, A'_4\}$. Since $a_1 a_3 \in E(G'')$, wlog. $A'_1 \cup A'_2 = a_1 \cup a_3$ and $A'_3 \cup A'_4 = a_2 \cup a_4$. By Proposition 3(a), $d_{G''}(s, A'_i) \geq 1$ so $d_G(s, A'_i) \geq 2$ $i \in \{1, 2\}$. By (1.28) and (1.29) for \mathcal{A} in G' , there exist $1 \leq l \leq r$ and $j \in \{1, 2\}$ such that for every edge $sd \in E(G')$ with $d \in A_j \cup A_{j+2}$, $c(sd) = l$. Then there exist $sd_1, sd_2 \in E(G'')$ with $d_1 \in A_j, d_2 \in A_{j+2}$ and $c(sd_1) = c(sd_2) = l$. This contradicts (1.29) for \mathcal{A}' .

Case b: Either we reduce this case to a case already seen or we show that the following edge pair e', f' will do. Let $g := sa_3$. If $c(g) \neq c(e)$, then let $e' := e$ and $f' := g$, otherwise, let $e' := g$ and $f' := f$. Since $c(e') = c(e)$,

G'' satisfies (1.37). By Propositions 1(a) and 3(c), $\{e', f'\}$ is admissible in G , so allowed in G and (1.34) is satisfied. If G'' satisfies (1.36), then we are done. Otherwise, either G'' contains a C_6 -obstacle, and then, by Lemma 5, we are done, or G'' contains a C_4 -obstacle $\mathcal{A}' := \{A'_1, A'_2, A'_3, A'_4\}$. Wlog. $x', y' \in A'_1$, otherwise restarting the proof by e' and f' we are in Case a. Then, by (1.26) and (1.81), $|A'_j| = 1 \forall 2 \leq j \leq 4$ and $d_G(A'_1) = k + 2$. By Proposition 2(b), $G[A'_1]$ is connected, so, by (1.27), wlog. $V - a_4 \subseteq A'_1 \cup A_1$. First suppose that $A'_1 \cup A_1 = V$. Then, by Propositions 3(b) applied for G' , $d_{G'}(A'_1) = k + 2$ and $k = d_{G'}(V - A'_1)$. The common edge of $\{e, f\}$ and $\{e', f'\}$ enters $A_1 \cap A'_1$, so $d_{G'}(V - A'_1) = d_G(V - A'_1)$ thus, by (1.81), $1 = |V - A'_1| (= 3)$, contradiction, so $A'_1 \cup A_1 = V - a_4$. Then, by Proposition 3(b) again applied for G' , $d_{G'}(A'_1 \cup A_1) \geq k + 2$. Then, by (1.2) and (1.34), $d_G(A'_1 \cap A_1) = k$ thus, by (1.81), $|A'_1 \cap A_1| = 1$, say $A'_1 \cap A_1 = a_1$. Then it follows that $|V| = 6$, say $V = \{a_1, a_2, \dots, a_6\}$. Note that $d_G(a_i) = k \ 1 \leq i \leq 6$. By Proposition 3(a) for \mathcal{A} and for \mathcal{A}' , $1 \leq d_G(sa_i) \ 1 \leq i \leq 6$ so $6 \leq d_G(s)$. The following claim provides a contradiction.

Claim. $\{a_1, a_2, \dots, a_6\}$ forms a C_6 -obstacle in G .

Proof. We start with some structural observations.

Proposition 4. (a) $A_1 = \{a_1, a_5, a_6\}$, $A_i = a_i \ 2 \leq i \leq 4$, $A'_1 = \{a_1, a_2, a_3\}$, wlog. $A'_2 = a_6$, $A'_3 = a_5$, $A'_4 = a_4$, (b) $d_G(a_1, a_2) = d_G(a_2, a_3) = d_G(a_1, a_6) = d_G(a_5, a_6) = \frac{k-1}{2}$, (c) $\{x, y\} = \{a_1, a_5\}$.

Proof. We know that $A'_1 = \{a_1, a_2, a_3\}$ and $A_1 = \{a_1, a_5, a_6\}$. Then, by (1.27) for \mathcal{A} , $d_G(a_1, a_3) = 0$ so, by Proposition 2(c), $d_G(a_1, a_2) = d_G(a_2, a_3) = \frac{k-1}{2}$. Wlog. $A'_2 = a_6$. Suppose that $A'_4 = a_5$. Then, by (1.27) for \mathcal{A}' , $d_G(a_5, a_6) = 0$ so, by Proposition 2(c), $d_G(a_1, a_5) = d_G(a_1, a_6) = \frac{k-1}{2}$. Then $k = d_G(a_1) \geq d_G(a_1, a_2) + d_G(a_1, a_5) + d_G(a_1, a_6) + d_G(a_1, s) \geq 3\frac{k-1}{2} + 1$, that is $k \leq 1$, contradiction. Thus $A'_4 = a_4$ and $A'_3 = a_5$, that is, (a) is satisfied. Then, by (1.27) for \mathcal{A}' , $d_G(a_1, a_5) = 0$ so, by Proposition 2(c), $d_G(a_1, a_6) = d_G(a_5, a_6) = \frac{k-1}{2}$ and (b) is satisfied. By definition, $\{x, y\} \cap \{x', y'\} = a_1$, so $a_1 \in \{x, y\}$. Suppose $a_5 \notin \{x, y\}$. Then $\{x, y\} = \{a_1, a_6\}$. By (1.2), (1.30) and (b), $d_G(\{a_1, a_6\}) = d_G(a_1) + d_G(a_6) - 2d_G(a_1, a_6) = k + k - (k - 1) = k + 1$, hence, by Proposition 1(a), $\{sx, sy\}$ is not admissible in G , contradiction, thus (c) is satisfied. \square

By (1.29) for \mathcal{A}' , $c(sa_2) \neq c(sa_4)$ so $\delta_{G'}(A_1 \cup A_3) \cap \delta_{G'}(s) = P'_l$ in (1.29) for \mathcal{A} for some l with $|P_l| \geq |P'_l| = \frac{d_{G'}(s)}{2} = \frac{d_G(s)}{2} - 1$. By (1.29) for \mathcal{A} , $c(sa_6) \neq c(sa_4)$ so $\delta_{G''}(A'_1 \cup A'_3) \cap \delta_{G''}(s) = P_{l'}$ in (1.29) for \mathcal{A}' for some l' with $|P_{l'}| \geq |P'_{l'}| = \frac{d_{G''}(s)}{2} = \frac{d_G(s)}{2} - 1$. In particular, $c(sa_2) = c(sa_5) = l'$. By (1.29) for \mathcal{A} , $l = c(sa_3) \neq c(sa_2) = l'$ thus, by Proposition 4(c), $e = e' = sa_1$, $f = sa_5$, $f' = sa_3$. Since $\{e, f\}$ and $\{e', f'\}$ are allowed, $l \neq 1 \neq l'$. Then, by the maximality of P_1 , $|P_1| \geq |P_l| \geq \frac{d_G(s)}{2} - 1$. $d_G(s) \geq |P_1| + |P_l| + |P_{l'}| \geq 3(\frac{d_G(s)}{2} - 1)$, that is, $d_G(s) \leq 6$. Then $d_G(s) = 6$ and $|P_1| = |P_l| = |P_{l'}| = 2$,

namely $P_1 = \{sa_1, sa_4\}$, $P_l = \{sa_3, sa_6\}$, $P_{l'} = \{sa_2, sa_5\}$, so (1.32) and (1.33) are satisfied. We have already seen that (1.30) is satisfied. By (1.26) and (1.27) for \mathcal{A}' and for \mathcal{A} , Proposition 4(b) and (1.32), $d_G(a_5, a_4) = \frac{k-1}{2} = d_G(a_3, a_4)$. Then, by Proposition 4(b), (1.31) is satisfied. This completes the proof of the claim and also of Lemma 6. \square

By Lemmas 4, there exists an allowed pair, and then, by Lemmas 5 and 6, there exists one that does not create a C_4 - or C_6 -obstacle, so by induction, there exists a complete allowed splitting off and Theorem 15 is proved. $\square \square \square$

Acknowledgement. I thank T. Király for the fruitful discussions on this topic.

References

1. J. Bang-Jensen, A. Frank, B. Jackson, Preserving and increasing local edge-connectivity in mixed graphs, *SIAM J. Discrete Math.* Vol. 8 No. 2 (1995) 155-178.
2. J. Bang-Jensen, H. Gabow, T. Jordán, Z. Szigeti, Edge-connectivity augmentation with partition constraints, *SIAM J. Discrete Math.* Vol. 12 No. 2 (1999) 160-207.
3. J. Bang-Jensen, B. Jackson, Augmenting hypergraphs by edges of size two, *Math. Program.* Vol. 84 No. 3 (1999) 467-481.
4. J. Bang-Jensen, T. Jordán, Edge-connectivity augmentation preserving simplicity, *SIAM J. Discrete Math.* Vol. 11 No. 4 (1998) 603-623.
5. J. Bang-Jensen, T. Jordán, Splitting off edges within a specified subset preserving the edge-connectivity of the graph, *J. of Algorithms* 37 (2000) 326-343.
6. A. Benczúr, A. Frank, Covering symmetric supermodular functions by graphs, *Math. Program.* Vol. 84 No. 3 (1999) 483-503.
7. A. Bernáth, T. Király, personal communication
8. G. R. Cai, Y. G. Sun, The minimum augmentation of any graph to k -edge-connected graphs, *Networks*, 19 (1989) 151-172.
9. E. Cheng, T. Jordán, Successive edge-connectivity augmentation problems, *Math. Program.* Vol. 84 No. 3 (1999) 577-593.
10. B. Cosh, Vertex Splitting and Connectivity Augmentation in Hypergraphs, Ph.D. thesis, University of London (2000)
11. B. Cosh, B. Jackson, Z. Király, Local connectivity augmentation in hypergraphs is NP-complete, submitted
12. K. P. Eswaran, E. Tarjan, Augmentation problems, *SIAM J. Comput.*, 5 (1976) 653-665.
13. T. Fleiner, T. Jordán, Covering and structure of crossing families, *Math. Program.* Vol. 84 No. 3 (1999) 505-518.
14. B. Fleiner, Detachments of vertices of graphs preserving edge-connectivity, *SIAM J. Discrete Math.* Vol. 18 No. 3 (2005) 581-591.
15. A. Frank, Augmenting graphs to meet edge-connectivity requirements, *SIAM J. Discrete Math.* Vol. 5 No. 1 (1992) 22-53.
16. A. Frank, On a theorem of Mader, *Discrete Math.* **101** (1992) 49-57.

17. A. Frank, oral communication
18. R. Grappe, Z. Szigeti, Covering semi-monotone symmetric functions, *Discrete Appl. Math.* 156 (2008) 138-144.
19. T. Ishii, M. Hagiwara, Minimum augmentation of local edge-connectivity between vertices and vertex subsets in undirected graphs, *Disc. Appl. Math.* 154 (2006) 2307-2329.
20. T. Ishii, Minimum augmentation of edge-connectivity with monotone requirements in undirected graphs, *Proc. of the 13th Australasian Symp. on Theory of Comp.*, (2007) 91-100.
21. B. Jackson, Some remarks on connectivity, vertex splitting and orientation in digraphs, *J. Graph Theory*, 12, (1988) 429-436.
22. T. Jordán, Two NP-Complete Augmentation Problems, IMADA, Odense Universitet, Preprints No. 8/1997.
23. T. Jordán, Constrained edge-splitting problems, *SIAM J. Discrete Math.* Vol. 17 No. 1 (2003) 88-102.
24. T. Jordán, Z. Szigeti, Detachments preserving local edge-connectivity of graphs, *SIAM J. Discrete Math.* Vol. 17 No. 1 (2003) 72-87.
25. T. Király, Merging hyperedges to meet edge-connectivity requirements, Egres Technical Reports, TR-2005-08 <http://www.cs.elte.hu/egres/>, 2004.
26. T. Király, Covering symmetric supermodular functions by uniform hyperedges, *J. Combin. Theory Ser. B* 91 (2004) 185-200.
27. Z. Király, oral communication
28. L. Lovász, Combinatorial problems and exercises, North-Holland, Amsterdam, (1979).
29. W. Mader, A reduction method for edge-connectivity in graphs, *Ann. Discrete Math.* 3 (1978) 145-164.
30. Z. Nutov, Approximating connectivity augmentation problems, In SODA 2005, 176-185.
31. Z. Szigeti, Hypergraph connectivity augmentation, *Math. Program.* Vol. 84 No. 3 (1999) 519-527.
32. Z. Szigeti, Edge-connectivity augmentation of graphs over symmetric parity families, to appear in *Discrete Math.*
33. Z. Szigeti, Edge-splittings preserving edge-connectivity of graphs, *Discrete Applied Mathematics*, 156 (2008) 1011-1018.
34. Z. Szigeti, A short proof on the local detachment theorem, Egres Technical Reports, TR-2004-09 <http://www.cs.elte.hu/egres/>, 2004.
35. Z. Szigeti, On partition constrained splitting off, Egres Technical Reports, TR-2004-08 <http://www.cs.elte.hu/egres/>, 2004.
36. T. Watanabe, A. Nakamura, Edge-connectivity augmentation problems, *Comp. System Sci.* 35 (1987) 96-144.