

# Edge-splittings preserving local edge-connectivity of graphs <sup>★</sup>

Zoltán Szigeti <sup>1</sup>

*Laboratoire Leibniz-IMAG  
46 avenue Félix Viallet  
Grenoble, France.*

---

## Abstract

Let  $G = (V + s, E)$  be a 2-edge-connected graph with a designated vertex  $s$ . A pair of edges  $rs, st$  is called admissible if splitting off these edges (replacing  $rs$  and  $st$  by  $rt$ ) preserves the local edge-connectivity (the maximum number of pairwise edge disjoint paths) between each pair of vertices in  $V$ . The operation splitting off is very useful in graph theory, it is especially powerful in the solution of edge-connectivity augmentation problems as it was shown by Frank [4]. Mader [7] proved that if  $d(s) \neq 3$  then there exists an admissible pair incident to  $s$ . We generalize this result by showing that if  $d(s) \geq 4$  then there exists an edge incident to  $s$  that belongs to at least  $\lfloor d(s)/3 \rfloor$  admissible pairs. An infinite family of graphs shows that this bound is best possible. We also refine a result of Frank [5] by describing the structure of the graph if an edge incident to  $s$  belongs to no admissible pairs. This provides a new proof for Mader's theorem.

*Keywords:* local edge-connectivity, splitting off, edge-connectivity augmentation

---

<sup>★</sup> This work was done while the author was visiting the Egerváry Research Group (EGRES), Department of Operations Research, Eötvös University, Budapest, Hungary.

<sup>★★</sup>An extended abstract of this paper was presented at GRACO2005 (2nd Brazilian Symposium on Graphs, Algorithms, and Combinatorics) and appeared, under a different title, in *Electronic Notes in Discrete Mathematics* 19 (2005) 57–61.

<sup>1</sup> Email: [Zoltan.Szigeti@imag.fr](mailto:Zoltan.Szigeti@imag.fr)

## 1 Introduction

In this paper,  $G = (V + s, E)$  denotes a 2-edge-connected graph,  $s$  being a vertex not in  $V$ . (It would be enough to suppose that no cut edge is incident to  $s$  but for the sake of simplicity we suppose that  $G$  contains no cut edge at all.)

For two vertices  $u, v \in V$ , the **local edge-connectivity**,  $\lambda_G(u, v)$ , between  $u$  and  $v$  is the maximum number of edge disjoint paths between  $u$  and  $v$ . If  $\lambda_G(u, v) \geq k$  for all pairs  $u, v \in V$ , then  $G$  is called  **$k$ -edge-connected in  $V$** .

The operation **splitting off** is defined as follows: two edges  $rs$  and  $st$  are replaced by a new edge  $rt$ . The graph obtained from  $G$  by splitting off a pair of edges  $rs, st$  is denoted by  $G_{rt}$ . A pair of edges  $rs, st$  is called  **$k$ -admissible** if  $G_{rt}$  is  $k$ -edge-connected in  $V$ . The pair of edges  $rs, st$  is called **admissible** if  $\lambda_{G_{rt}}(u, v) \geq \lambda_G(u, v)$  for all pairs  $u, v \in V$ . An edge incident to  $s$  is called **admissible** if it belongs to an admissible pair, otherwise it is called **non-admissible**.

The first splitting off result is due to Lovász [6].

**Theorem 1.1** *If  $G = (V + s, E)$  is  $k$ -edge-connected in  $V$  for some  $k \geq 2$  and  $d(s)$  is even then each edge incident to  $s$  belongs to a  $k$ -admissible pair.*

Cai and Sun [3] showed how to apply this result to solve the following global edge-connectivity augmentation problem: Given a graph  $H$  and an edge-connectivity requirement  $k \in \mathbb{Z}_+$ , find the minimum number of new edges whose addition makes the graph  $k$ -edge-connected.

Theorem 1.1 was extended in Bang-Jensen et al. [1].

**Theorem 1.2** *If  $G = (V + s, E)$  is  $k$ -edge-connected in  $V$  for some  $k \geq 2$  and  $d(s)$  is even then each edge incident to  $s$  belongs to at least  $d(s)/2$  (resp.  $d(s)/2 - 1$ )  $k$ -admissible pairs if  $k$  is even (resp. odd).*

In [1], we applied Theorem 1.2 to solve the global edge-connectivity augmentation problem in bipartite graphs: Given a connected bipartite graph  $H$  and an edge-connectivity requirement  $k \in \mathbb{Z}_+$ , what is the minimum number of new edges whose addition results in a bipartite  $k$ -edge-connected graph.

It is easy to construct examples to show that the bounds of Theorem 1.2 are best-possible.

Mader [7] generalized Theorem 1.1 on local edge-connectivity.

**Theorem 1.3** *If  $G = (V + s, E)$  is 2-edge-connected and  $d(s) \neq 3$  then there exists an admissible pair incident to  $s$ .*

Applying this result, Frank [5] solved the local edge-connectivity augmentation problem: Given a graph  $H = (V, E)$  and a requirement function  $r : V \times V \rightarrow \mathbb{Z}_+$ , find the minimum number of new edges  $F$  such that  $\lambda_{H+F}(u, v) \geq r(u, v)$  for all pairs  $u, v \in V$ .

The main contribution of the present paper is the following strengthening of Theorem 1.3. It can be considered as the counterpart of Theorem 1.2 for local edge-connectivity.

**Theorem 1.4** *If  $G = (V + s, E)$  is a 2-edge-connected graph and  $d(s) \geq 4$  then there is an edge  $sr$  that belongs to at least  $\lfloor d(s)/3 \rfloor$  admissible pairs incident to  $s$ .*

We present, in Section 3, an infinite family of graphs showing that our bound is best possible.

Theorem 1.3 implies that at most three edges incident to  $s$  are non-admissible. Frank [5] provided a slight generalization of this result.

**Theorem 1.5** *If  $G = (V + s, E)$  is 2-edge-connected and  $d(s) \neq 3$  then at most one edge incident to  $s$  belongs to no admissible pair.*

We refine this result by describing the structure of the graph if it contains a non-admissible edge incident to  $s$ . (For definitions, see Section 2.)

**Theorem 1.6** *Let  $st$  be an edge of a 2-edge-connected graph  $G = (V + s, E)$ . The following are equivalent.*

- (a) *The edge  $st$  is non-admissible,*
- (b) *there exist two dangerous sets  $M_1$  and  $M_2$  such that  $t \in M_1 \cap M_2$  and  $M_1 \cup M_2$  contains all the neighbours of  $s$ ,*
- (c) *the degree  $d(s)$  of  $s$  is odd and there exist two disjoint tight sets  $C_1$  and  $C_2$  in  $V - t$  such that  $d(s, C_1) = d(s, C_2) = (d(s) - 1)/2$ .*

As an application of Theorem 1.6 we present the following result.

**Theorem 1.7** *Let  $G = (V + s, E)$  be a 2-edge-connected graph with  $d(s) \neq 3$ . If an edge  $st$  is non-admissible then each edge  $sr \neq st$  belongs to exactly  $(d(s) - 1)/2$  admissible pairs.*

The proofs of Theorems 1.6 and 1.7, given in Sections 4 and 5, together provides a new proof of Theorem 1.5 and hence of Theorem 1.3.

We mention a related interesting result of Bang-Jensen and Jordán.

**Theorem 1.8** [2] *Let  $G = (V + s, E)$  be a 2-edge-connected graph. Then, for every edge  $st$ , the number of edges  $rs$  for which the pair of edges  $rs, st$  is non-admissible is at most  $2k^2 - 2k$ , where  $k = \max\{\lambda_G(u, v) : u, v \in V\}$ .*

## 2 Notation and preliminary results

Let  $G = (V + s, E)$  be a graph, with  $s$  a vertex not in  $V$ . Let  $\Gamma(s)$  denote the set of neighbours of  $s$ . We use the notation  $\subset$  for proper subset. For a set  $T \subset V, T \neq \emptyset$  we denote the graph obtained from  $G$  by contracting  $T$  into one vertex  $v_T$  by  $G/T$ .

Let  $X, Y \subseteq V + s$ . Let  $d(X, Y)$  denote the number of edges between  $X - Y$  and  $Y - X$ . Let  $\bar{d}(X, Y)$  denote the number of edges between  $X \cap Y$  and  $V + s - (X \cup Y)$ . We define the degree of the set  $X$  by  $d(X) = d(X, V + s - X)$ . The degree function satisfies the following two well-known equalities.

- (1)  $d(X) + d(Y) = d(X \cap Y) + d(X \cup Y) + 2d(X, Y)$ ,
- (2)  $d(X) + d(Y) = d(X - Y) + d(Y - X) + 2\bar{d}(X, Y)$ .

Observe that, by Menger's theorem,  $\lambda_G(x, y) = \lambda(x, y) = \min\{d(Z) : Z \subset V + s, x \in Z, y \notin Z\}$  for all  $x, y \in V$ . We define the function  $R(X)$  as follows:  $R(\emptyset) = R(V) = 0$  and for a set  $X \subset V, X \neq \emptyset$ , let

$$R(X) = \max\{\lambda_G(x, y) : x \in X, y \in V - X\}.$$

Observe that the function  $R(X)$  satisfies (3) and (4) for  $X, Y \subset V$ .

- (3)  $R(X) = R(V - X)$ ,
- (4)  $R((X - Y) \cup (Y - X)) \leq \max\{R(X - Y), R(Y - X)\}$ .

The following property of  $R(X)$  can be found in [4, Proposition 5.4]: for  $X, Y \subset V$ , at least one of (5) and (6) hold. If  $X \cup Y = V$  then (6) holds.

- (5)  $R(X) + R(Y) \leq R(X \cap Y) + R(X \cup Y)$ ,
- (6)  $R(X) + R(Y) \leq R(X - Y) + R(Y - X)$ .

Finally, we define the function

$$h(X) := d(X) - R(X).$$

Note that the function  $h(X)$  satisfies (7) and (8) for  $X, Y \subset V$ .

- (7)  $h(X) \geq 0$ ,
- (8)  $h(X) = h(V - X) + 2d(s, X) - d(s)$ .

The properties above imply

**Proposition 2.1** For  $X, Y \subset V$ , at least one of (9) and (10) hold. If  $X \cup Y = V$  then (10) holds.

$$(9) \quad h(X) + h(Y) \geq h(X \cap Y) + h(X \cup Y) + 2d(X, Y),$$

$$(10) \quad h(X) + h(Y) \geq h(X - Y) + h(Y - X) + 2\bar{d}(X, Y).$$

A set  $\emptyset \neq X \subset V$  is called **tight** if  $h(X) = 0$  and it is called **dangerous** if  $h(X) \leq 1$ . Note that tight and dangerous sets are, by definition, subsets of  $V$ .

The following claim is due to Mader.

**Claim 2.2** Let  $T$  be a tight set in a graph  $G = (V + s, E)$  and  $G' := G/T$ .

(a) [7, Lemma 3] If a pair of edges  $e', f'$  incident to  $s$  is admissible in  $G'$  then the corresponding pair of edges  $e, f$  is admissible in  $G$ .

(b) [7, Lemma 4] If  $X' \subseteq V(G') - s$  then  $h_{G'}(X') = h_G(X)$ , where  $X = X' - v_T \cup T$  if  $v_T \in X'$  and  $X = X'$  otherwise.

The reduction method of Claim 2.2 will be applied in our proofs and hence we will be able to assume that

(11) every tight set is a singleton.

We need the following claims.

**Claim 2.3** [5, Claim 3.1] A pair of edges  $us, sv$  of a graph  $G = (V + s, E)$  is admissible if and only if there is no dangerous set  $M$  with  $u, v \in M$ .

**Claim 2.4** [5, Claim 4.1] Let  $G = (V + s, E)$  be a graph and  $t \in \Gamma(s)$  be a vertex of minimum degree. Suppose that (11) holds. If a set  $M \subseteq V$  contains  $t$  and  $|\Gamma(s) \cap M| \geq 2$ , then  $R(M - t) \geq R(M)$ .

**Claim 2.5** Let  $G = (V + s, E)$  be a 2-edge-connected graph. If  $M$  is a dangerous set then

(a)  $d(s, M) \leq (d(s) + 1)/2$ , with equality only if  $V - M$  is tight, and

(b) [2, in Lemma 5.4]  $d(X, M - X) \geq 1$  for every  $\emptyset \neq X \subset M$ .

**Proof.** (a) By (8), since  $M$  is dangerous and by applying (7) for  $V - M$ ,  $d(s, M) = (d(s) + h(M) - h(V - M))/2 \leq (d(s) + 1)/2$  and (a) follows.  $\square$

We close this section with a technical lemma.

**Lemma 2.6** Let  $G = (V + s, E)$  be a 2-edge-connected graph,  $st \in E$  and  $S \subseteq V$ . Let  $\mathcal{M}$  be a minimum collection of dangerous sets such that  $t \in \bigcap \mathcal{M}$  and  $S \subseteq \bigcup \mathcal{M}$ . If  $|\mathcal{M}| \geq 3$ , (11) holds and  $M_i, M_j \in \mathcal{M}$ , then

(a) (10) does not apply for  $M_i$  and  $M_j$ , and

(b)  $M_i \cap M_j = t$ .

**Proof.** (a) Suppose that (10) applies for  $M_i$  and  $M_j$ . Then, by  $1 \geq h(M_i)$  and  $1 \geq h(M_j)$ , we have  $h(M_i - M_j) = 0$  and  $h(M_j - M_i) = 0$  (so by (11),  $M_i - M_j = r_i$  and  $M_j - M_i = r_j$  for some vertices  $r_i, r_j \in V$ ) and  $\bar{d}(M_i, M_j) = 1$ . Let  $M_k \in \mathcal{M} - \{M_i, M_j\}$  and  $X = M_i \cap M_j \cap M_k$ . Note that  $t \in X$  so  $X \neq \emptyset$ . By the minimality of  $\mathcal{M}$ ,  $M_k - X \neq \emptyset$ . Then, by Claim 2.5(b) and since  $st$  enters  $M_i \cap M_j$ , we have  $1 \leq d(X, M_k - X) \leq d(M_i \cap M_j, M_k - (M_i \cap M_j)) \leq \bar{d}(M_i, M_j) - d(M_i \cap M_j, s) \leq 1 - 1 = 0$ , a contradiction.

(b) By Proposition 2.1 and (a), (9) applies for  $M_i$  and  $M_j$ . Then, since  $1 \geq h(M_i)$ ,  $1 \geq h(M_j)$ , and by the minimality of  $\mathcal{M}$ ,  $h(M_i \cup M_j) \geq 2$  (otherwise we could replace  $M_i$  and  $M_j$  by  $M_i \cup M_j$ ), we have  $h(M_i \cap M_j) = 0$  and hence, by (11) and  $t \in M_i \cap M_j$ , (b) is satisfied.  $\square$

### 3 Proof of Theorem 1.4

The proof is similar to the proof of Theorem 1.3 given by Frank in [5].

**Proof.** We prove the theorem by induction on  $|V|$ . We may assume, by Claim 2.2(a), that (11) is satisfied. Let  $t$  be a neighbour of  $s$  of minimum degree. Let  $S$  be the set of neighbours  $r$  of  $s$  such that  $r = t$  or the pair of edges  $rs, st$  is not admissible. By Claim 2.3, there is a minimum collection  $\mathcal{M}$  of dangerous sets such that  $t \in \bigcap \mathcal{M}$  and  $S \subseteq \bigcup \mathcal{M}$ . Suppose that  $st$  belongs to less than  $\lfloor d(s)/3 \rfloor$  admissible pairs (otherwise, we are done). Then

$$(12) \quad d(s, \bigcup \mathcal{M}) \geq d(s, S) > d(s) - \lfloor d(s)/3 \rfloor = \lceil 2d(s)/3 \rceil.$$

By Claim 2.5(a) and (12), for  $M_i \in \mathcal{M}$ ,  $d(s, M_i) \leq (d(s) + 1)/2 < \lceil 2d(s)/3 \rceil < d(s, \bigcup \mathcal{M})$  and hence  $|\mathcal{M}| \geq 2$ . Let  $M_1, M_2 \in \mathcal{M}$ . By the minimality of  $\mathcal{M}$ , each  $M_i \in \mathcal{M}$  contains a neighbour  $r_i \neq t$  of  $s$  that belongs to no other  $M_j \in \mathcal{M}$ . Let us choose such a vertex  $r_i$  for each  $M_i \in \mathcal{M}$ .

**Claim 3.1**  $\mathcal{M} = \{M_1, M_2\}$ .

**Proof.** For  $i = 1, 2$ ,  $M_i$  contains  $t$  and  $r_i$ , so  $|\Gamma(s) \cap M_i| \geq 2$ . Then, by Claim 2.4,  $R(M_1 - t) \geq R(M_1)$  and  $R(M_2 - t) \geq R(M_2)$ . Suppose that  $|\mathcal{M}| \geq 3$ . Then, by Lemma 2.6(b),  $M_1 \cap M_2 = t$ , thus  $M_1$  and  $M_2$  satisfy (6) and hence (10), a contradiction by Lemma 2.6(a).  $\square$

**Claim 3.2** (10) applies for  $M_1$  and  $M_2$ .

**Proof.** Suppose that (10) does not hold for  $M_1$  and  $M_2$ . Then, by Proposition 2.1,  $M_1 \cup M_2 \neq V$  and (9) applies for  $M_1$  and  $M_2$ . By (8), (7), Claim 3.1, (12) and  $d(s) \geq 4$ ,  $h(M_1 \cup M_2) \geq 2d(s, M_1 \cup M_2) - d(s) = 2d(s, \bigcup \mathcal{M}) - d(s) >$

$2\lceil 2d(s)/3 \rceil - d(s) \geq 2$ . It follows, by  $1 \geq h(M_1), 1 \geq h(M_2)$ , (9) and (7), that  $1+1 \geq h(M_1) + h(M_2) \geq h(M_1 \cap M_2) + h(M_1 \cup M_2) > 0+2$ , a contradiction.  $\square$

**Claim 3.3**  $d(s, r_1) + d(s, r_2) \geq \lceil 2d(s)/3 \rceil$ .

By  $1 \geq h(M_1), 1 \geq h(M_2)$ , Claim 3.2, (7),  $st \in E$  and  $t \in M_1 \cap M_2$ , we have  $1+1 \geq h(M_1) + h(M_2) \geq h(M_1 - M_2) + h(M_2 - M_1) + 2\bar{d}(M_1, M_2) \geq 0+0+2d(s, M_1 \cap M_2) \geq 2$ , so  $h(M_1 - M_2) = 0 = h(M_2 - M_1)$  and  $d(s, M_1 \cap M_2) = 1$ . It follows, by  $r_1 \in M_1 - M_2, r_2 \in M_2 - M_1$  and (11), that  $M_1 - M_2 = r_1$  and  $M_2 - M_1 = r_2$ . Then, by Claim 3.1 and (12),  $d(s, r_1) + d(s, r_2) = d(s, M_1 \cup M_2) - d(s, M_1 \cap M_2) = d(s, \bigcup \mathcal{M}) - 1 \geq \lceil 2d(s)/3 \rceil$ .  $\square$

Let  $e_i$  be any edge connecting  $s$  and  $r_i$  for  $1 \leq i \leq 2$ .

**Claim 3.4** *The pair of edges  $e_1, e_2$  is admissible.*

**Proof.** Otherwise, by Claim 2.3, there is a dangerous set  $X$  with  $r_1, r_2 \in X$ , and then, by (8), (7), Claim 3.3 and  $d(s) \geq 4$ , we have  $1 \geq h(X) \geq 2d(s, X) - d(s) \geq 2\lceil 2d(s)/3 \rceil - d(s) \geq 2$ , a contradiction.  $\square$

By Claim 3.3, we may assume without loss of generality that  $d(s, r_1) \geq \lceil d(s)/3 \rceil \geq \lfloor d(s)/3 \rfloor$ . Then, by Claim 3.4,  $e_2$  belongs to at least  $\lfloor d(s)/3 \rfloor$  admissible pairs and the proof of Theorem 1.4 is complete.  $\square$

**Examples:** There exists an infinite class of graphs in which each edge incident to  $s$  belongs to exactly  $\lfloor d(s)/3 \rfloor$  admissible pairs. See Figure 1. We mention that it is not true in general, even if we suppose that the degree of  $s$  is even, that each edge incident to  $s$  belongs to many admissible pairs. In Figure 2, the edge  $ws$  belongs to the unique admissible pair of edges  $ws, sz$ .

## 4 Proof of Theorem 1.6

**Proof.** We consider first the most complicated part, we prove that (a) implies (b) by induction on  $|V|$ .

**Claim 4.1** *We may assume that (11) is satisfied.*

**Proof.** Suppose that there exists a tight set  $T$  with  $|T| > 1$ . Let  $G' = G/T$ . By Claim 2.2(a),  $st$  belongs to no admissible pair in  $G'$ ,  $G'$  is 2-edge-connected and  $|V(G')| < |V|$ , hence, by induction, (b) is true for  $G'$  and then, by Claim 2.2 (b), it is also true for  $G$ .  $\square$

The edge  $st$  belongs to no admissible pair, thus, by Claim 2.3, there is a minimum collection  $\mathcal{M}$  of dangerous sets such that  $t \in \bigcap \mathcal{M}$  and  $\Gamma(s) \subseteq \bigcup \mathcal{M}$ . By the minimality of  $\mathcal{M}$ , each  $M_i \in \mathcal{M}$  contains a neighbour  $r_i \neq t$




Fig. 1. Each edge incident to  $s$  belongs to exactly  $\lfloor d(s)/3 \rfloor$  admissible pairs.




Fig. 2. The degree  $d(s)$  of  $s$  is even and the edge  $ws$  belongs to a unique admissible pair  $ws, sz$ .

of  $s$  that belongs to no other  $M_j \in \mathcal{M}$ . Let us choose such a vertex  $r_i$  for each  $M_i \in \mathcal{M}$ . By Claim 2.5(a),  $d(s) \geq 2$  and  $\Gamma(s) \subseteq \bigcup \mathcal{M}$ , for  $M_i \in \mathcal{M}$ ,  $d(s, M_i) \leq (d(s) + 1)/2 < d(s) = d(s, \bigcup \mathcal{M})$  and hence  $|\mathcal{M}| \geq 2$ .

Suppose that  $|\mathcal{M}| \geq 3$ . We shall find a contradiction showing that this case can not happen and hence  $|\mathcal{M}| = 2$ . By Lemma 2.6(b), for all  $M_i, M_j \in \mathcal{M}$ ,  $M_i - M_j = M_i - t$ . Let  $T = V - \bigcup \mathcal{M}$ . Note that  $d(s, T) = 0$ .

**Claim 4.2** *If  $R(M_1) = \lambda(a, b)$  with  $a \in M_1$  and  $b \in T$ , then for some  $M_k \in$*



$\mathcal{M} - M_1$ ,  $R(M_k - t) > R(t)$ .

**Proof.** Note that  $d(s) \geq |\mathcal{M}| + 1$  and  $d(T) \geq \lambda(a, b) = R(M_1) \geq d(M_1) - 1$  because  $M_1$  is dangerous. By repeated applications of (1) we get

$$\begin{aligned} \sum_{M_j \in \mathcal{M}} (d(M_j) - d(t)) &\geq d(s \cup T) - d(t) \\ &= d(s) + d(T) - d(t) \\ &\geq (|\mathcal{M}| + 1) + (d(M_1) - 1) - d(t) \\ &> (|\mathcal{M}| - 1) + (d(M_1) - d(t)), \end{aligned}$$

so there exists  $M_k \in \mathcal{M} - M_1$  with  $d(M_k) - d(t) > 1$ . Then, since  $M_k$  is dangerous,  $R(M_k) \geq d(M_k) - 1 > d(t) \geq R(t)$  so, by (4),  $R(M_k - t) > R(t)$ .  $\square$

**Claim 4.3** *There exists  $M_i \in \mathcal{M}$  for which  $R(M_i - t) \geq R(t)$ .*

**Proof.** Let  $Y = \{y \in V - t : R(t) = \lambda(t, y)\}$ . By definition,  $Y \neq \emptyset$ . If there exists a vertex  $y \in M_i \cap Y$  for some  $M_i \in \mathcal{M}$ , then  $R(M_i - t) \geq \lambda(t, y) = R(t)$ . Thus we may suppose that  $Y \subseteq T$ . Let  $y \in Y$ . Then  $R(M_1) \geq \lambda(t, y) = R(t)$ . If  $R(M_1) = \lambda(t, y)$  then, by Claim 4.2,  $R(M_1 - t) > R(t)$ . Otherwise  $R(M_1) > R(t)$  so, by (4),  $R(M_1 - t) > R(t)$ .  $\square$

**Claim 4.4** *If  $M_j \in \mathcal{M} - M_i$ , then  $R(M_j - t) < R(M_j) \leq R(t)$ .*

**Proof.** Suppose that  $R(M_j - t) \geq R(M_j)$ . By Claim 4.3 and (4),  $R(M_i - t) \geq R(M_i)$ . So (6) and hence (10) applies for  $M_i$  and  $M_j$ , contradicting Lemma 2.6(a). By  $R(M_j - t) < R(M_j)$  and (4),  $R(M_j) \leq R(t)$ .  $\square$

**Claim 4.5** *If  $R(M_i) = \lambda(a, b)$  with  $a \in M_i$  and  $b \in V - M_i$ , then  $b \in T$ .*

**Proof.** Suppose that  $b \in M_j \in \mathcal{M} - M_i$ . Then,  $R(M_j - t) \geq \lambda(a, b) = R(M_i)$ . By Claims 4.4 and 4.3,  $R(M_j) \leq R(t) \leq R(M_i - t)$ . Thus (6) and hence (10) applies for  $M_i$  and  $M_j$ , a contradiction by Lemma 2.6(a).  $\square$

By Claims 4.3 and 4.4, there exists  $M_i \in \mathcal{M}$  such that  $R(M_j - t) < R(t)$  for all  $M_j \in \mathcal{M} - M_i$ . However, by Claim 4.5 and Claim 4.2, applied for  $M_1 = M_i$ ,  $R(M_j - t) > R(t)$  for some  $M_j \in \mathcal{M} - M_i$ . This contradiction completes the proof of (a) implies (b).

Obviously, (b) implies (a) by Claim 2.3.

We show now that (b) implies (c). Let  $C_1 = M_1 - M_2$  and  $C_2 = M_2 - M_1$ . Clearly,  $C_1 \cap C_2 = \emptyset$  and, by  $t \in M_1 \cap M_2$ , the sets  $C_1$  and  $C_2$  are in  $V - t$ .

**Claim 4.6**  *$d(s)$  is odd and  $d(s, C_1) = (d(s) - 1)/2 = d(s, C_2)$ .*

**Proof.** By (8),  $\Gamma(s) \subseteq M_1 \cup M_2$  and  $st \in E$ , we have  $2(d(s) + 1)/2 \geq d(s, M_1) + d(s, M_2) = d(s, M_1 \cup M_2) + d(s, M_1 \cap M_2) \geq d(s) + 1$ . It follows that  $d(s)$  is odd,  $d(s, M_i) = (d(s) + 1)/2$  and  $d(s, M_1 \cap M_2) = 1$ . Then  $d(s, C_i) = d(s, M_i) - d(s, M_1 \cap M_2) = (d(s) + 1)/2 - 1 = (d(s) - 1)/2$  for  $i = 1, 2$ .  $\square$

**Claim 4.7** (10) applies for  $M_1$  and  $M_2$ .

**Proof.** Suppose that (10) does not hold for  $M_1$  and  $M_2$ . Then, by Proposition 2.1,  $M_1 \cup M_2 \neq V$  and (9) applies for  $M_1$  and  $M_2$ , so, by  $1 \geq h(M_1), 1 \geq h(M_2)$  and (7), we have  $2 \geq h(M_1 \cup M_2)$ . It follows, by (8), (7) and  $\Gamma(s) \subseteq M_1 \cup M_2$ , that  $2 \geq h(M_1 \cup M_2) = h(V - (M_1 \cup M_2)) + 2d(s, M_1 \cup M_2) - d(s) \geq d(s)$ . However, since  $G$  is 2-edge-connected and  $d(s)$  is odd,  $d(s) \geq 3$ , a contradiction.  $\square$

Then, by  $1 \geq h(M_1), 1 \geq h(M_2)$ , (10),  $t \in M_1 \cap M_2$ , and  $st \in E$ , we get that  $h(C_1) = 0 = h(C_2)$ , that is  $C_1$  and  $C_2$  are tight sets. This completes the proof of (b) implies (c).

Finally, we show that (c) implies (b). Suppose that  $d(s)$  is odd and there exist two disjoint tight sets  $C_1, C_2 \subseteq V - t$  such that  $d(s, C_1) = (d(s) - 1)/2 = d(s, C_2)$ . Then, by (8),  $M_1 = V - C_1$  and  $M_2 = V - C_2$  are dangerous sets. Note that  $t \in M_1 \cap M_2$  and  $\Gamma(s) \subseteq M_1 \cup M_2$ .  $\square$

## 5 Proof of Theorem 1.7

**Proof.** By Theorem 1.6, there exist two dangerous sets  $M_1$  and  $M_2$  with  $t \in M_1 \cap M_2$  and  $\Gamma(s) \subseteq M_1 \cup M_2$ . It also follows from the proof above that  $d(s, M_1 \cap M_2) = 1$  and  $d(s, M_1) = d(s, M_2) = (d(s) + 1)/2$ . Let  $sr \neq st$  be an edge incident to  $s$ . Then, by Claim 2.3, the edge  $sr$  belongs to at most  $d(s) - (d(s) + 1)/2 = (d(s) - 1)/2$  admissible pairs. To finish the proof we show the following lemma.

**Lemma 5.1** *The edge  $sr$  belongs to at least  $(d(s) - 1)/2$  admissible pairs.*

**Proof.** We prove the lemma by induction on  $|V|$ . We may assume, by Claim 2.2(a), that (11) is satisfied. By Theorem 1.6,  $d(s)$  is odd and there exist two disjoint tight sets  $C_1, C_2 \subseteq V - t$  such that  $d(s, C_1) = d(s, C_2) = (d(s) - 1)/2$ . Then, by (11),  $C_1 = c_1$  and  $C_2 = c_2$  for some vertices  $c_1, c_2 \in V$ . Since  $sr \neq st$ , either  $r = c_1$  or  $r = c_2$ . The lemma follows from the following claim.

**Claim 5.2** *Let  $e_i$  be any edge connecting  $s$  and  $c_i$  for  $1 \leq i \leq 2$ . Then the pair of edges  $e_1, e_2$  is admissible.*

**Proof.** Otherwise, by Claim 2.3, there is a dangerous set  $X$  containing  $c_1$  and  $c_2$ . Then, by  $d(s, c_1) = d(s, c_2) = (d(s) - 1)/2$  and Claim 2.5(a),  $2(d(s) - 1)/2 \leq d(s, X) \leq (d(s) + 1)/2$ , that is  $d(s) \leq 3$ . However, since  $G$  is 2-edge-connected and  $d(s)$  is odd and  $\neq 3$ ,  $d(s) \geq 5$ , a contradiction.  $\square$

## 6 Open problems

For a summary on edge-connectivity augmentation problems in graphs we refer to [8]. We repeat one of the open problems proposed in [8], the problem of *local edge-connectivity augmentation in bipartite graphs*: given a connected bipartite graph  $H = (V, E)$  and a requirement function  $r : V \times V \rightarrow \mathbb{Z}_+$ , find the minimum number of new edges  $F$  such that  $\lambda_{H+F}(u, v) \geq r(u, v)$  for all pairs  $u, v \in V$  and  $H + F$  is a bipartite graph. Theorem 1.4 could help to solve this problem.

## References

- [1] J. Bang-Jensen, H. Gabow, T. Jordán, Z. Szigeti, Edge-connectivity augmentation with partition constraints, *SIAM J. Disc. Math.*, Vol. 12, No. 2 (1999), 160-207.
- [2] J. Bang-Jensen, T. Jordán, Edge-connectivity augmentation preserving simplicity, *SIAM J. Disc. Math.*, Vol. 11, No. 4 (1998), 603-623.
- [3] G. R. Cai, Y. G. Sun, The minimum augmentation of any graph to  $k$ -edge-connected graphs, *Networks*, 19 (1989), 151-172.
- [4] A. Frank, Augmenting graphs to meet edge-connectivity requirements, *SIAM J. Disc. Math.*, Vol. 5, No. 1 (1992), 22-53.
- [5] A. Frank, On a theorem of Mader, *Discrete Mathematics*, 101 (1992), 49-57.
- [6] L. Lovász, *Combinatorial problems and exercises*, North-Holland, Amsterdam, (1979).
- [7] W. Mader, A reduction method for edge-connectivity in graphs, *Ann. Discrete Math.*, 3 (1978), 145-164.
- [8] Z. Szigeti, Edge-connectivity augmentations of graphs and hypergraphs, Technical Report No. 04941 of the Research Institute for Discrete Mathematics, Bonn, 2004.
- [9] Z. Szigeti, On admissible edges, Technical Report 2004-07 of the Egerváry Research Group, Budapest, 2004, <http://www.cs.elte.hu/egres/>.